



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#72374 store at +4C

## CD20 (2H7) Mouse mAb (PerCP-Cy5.5<sup>®</sup> Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> FC-FP, FC-L	<b>Reactivity:</b> H	<b>Sensitivity:</b> Endogenous	<b>Source/Isotype:</b> Mouse IgG2b kappa	<b>UniProt ID:</b> #P11836	<b>Entrez-Gene Id:</b> 931
-------------------------------------	-------------------------	-----------------------------------	---	-------------------------------	-------------------------------

Product Usage Information	Application	Dilution
	Flow Cytometry (Fixed/Permeabilized)	1:20
	Flow Cytometry (Live)	1:20
<b>Storage</b>	Supplied in 10 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, 0.09% NaN <sub>3</sub> , 0.1% gelatin, pH 7.2. This product is stable for 6 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
<b>Specificity/Sensitivity</b>	CD20 (2H7) Mouse mAb (PerCP-Cy5.5 <sup>®</sup> Conjugate) recognizes endogenous levels of total CD20 protein. This antibody detects an epitope within the extracellular domain.	
<b>Source / Purification</b>	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.	
<b>Description</b>	This Cell Signaling Technology antibody is conjugated to PerCP-Cy5.5 <sup>®</sup> and tested in-house for direct flow cytometric analysis in human cells.	
<b>Background</b>	B-lymphocyte antigen CD20 (also known as MS4A1; Membrane-spanning 4-domains subfamily A member 1) is a cell surface phosphoprotein involved in the regulation of B cell activation and proliferation (1,2). It is commonly used as a marker to identify B cells and is expressed throughout B cell development, up until their differentiation into plasma cells. CD20 has no known ligand, and its expression and function are largely conserved between human and mouse (1-3). Evidence suggests that CD20 is necessary for store operated calcium (SOC) entry, which leads to elevated cytoplasmic calcium levels required for B cell activation (4,5). Anti-CD20 antibody immunotherapy depletes B cells by activation of the innate monocytic network and is a common treatment for B cell lymphomas, leukemias, and autoimmune diseases (6). The 2H7 antibody is widely used to identify both normal and malignant B cells (7).	
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Stashenko, P. et al. (1980) <i>J Immunol</i> 125, 1678-85.</li> <li>2. Tedder, T.F. et al. (1985) <i>J Immunol</i> 135, 973-9.</li> <li>3. Tedder, T.F. et al. (1988) <i>J Immunol</i> 141, 4388-94.</li> <li>4. Bubien, J.K. et al. (1993) <i>J Cell Biol</i> 121, 1121-32.</li> <li>5. Li, H. et al. (2003) <i>J Biol Chem</i> 278, 42427-34.</li> <li>6. Uchida, J. et al. (2004) <i>J Exp Med</i> 199, 1659-69.</li> <li>7. Liu, A.Y. et al. (1987) <i>J Immunol</i> 139, 3521-6.</li> </ol>	

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Applications Key</b>	<b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized) <b>FC-L:</b> Flow Cytometry (Live)
<b>Cross-Reactivity Key</b>	<b>H:</b> Human
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. Cy and CyDye are registered trademarks of GE Healthcare. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.