

5-Formylcytosine (5-fC) (D5D4K) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:
IF-IC, Dot Blot	All	Transfected Only	Rabbit IgG

Product Usage Information**Application**

Immunofluorescence (Immunocytochemistry)
DNA Dot Blot

Dilution

1:200
1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

5-Formylcytosine (5-fC) (D5D4K) Rabbit mAb detects 5-fC by IF in cells over-expressing the TET1 catalytic domain and by dot blot using double-stranded PCR fragments containing 5-fC. Many cells and tissues contain very low endogenous levels of 5-fC that may fall below the detection limits of this antibody. This antibody has been validated for specificity using ELISA and dot blot and shows high specificity for 5-fC.

Source / Purification

Monoclonal antibody is produced by immunizing animals with 5-formyl-2'-deoxycytosine.

Background

Methylation of DNA at cytosine residues is a heritable, epigenetic modification that is critical for proper regulation of gene expression, genomic imprinting, and mammalian development (1,2). 5-methylcytosine is a repressive epigenetic mark established *de novo* by two enzymes, DNMT3a and DNMT3b, and is maintained by DNMT1 (3, 4). 5-methylcytosine was originally thought to be passively depleted during DNA replication. However, subsequent studies have shown that Ten-Eleven Translocation (TET) proteins TET1, TET2, and TET3 can catalyze the oxidation of methylated cytosine to 5-hydroxymethylcytosine (5-hmC) (5). Additionally, TET proteins can further oxidize 5-hmC to form 5-formylcytosine (5-fC) and 5-carboxylcytosine (5-caC), both of which are excised by thymine-DNA glycosylase (TDG), effectively linking cytosine oxidation to the base excision repair pathway and supporting active cytosine demethylation (6,7). TET protein-mediated cytosine hydroxymethylation was initially demonstrated in mouse brain and embryonic stem cells (5, 8). Since then this modification has been discovered in many tissues, with the highest levels found in the brain (9). While 5-fC and 5-caC appear to be short-lived intermediate species, there is mounting evidence showing that 5-hmC is a distinct epigenetic mark with various unique functions (10,11). The modified base itself is stable *in vivo* and interacts with various readers including MeCP2 (11,12). The global level of 5-hmC increases during brain development, and 5-hmC is enriched at promoter regions and poised enhancers. Furthermore, there is an inverse correlation between levels of 5-hmC and histone H3K9 and H3K27 trimethylation, suggesting a role for 5-hmC in gene activation (12). Lower amounts of 5-hmC have been reported in various cancers including myeloid leukemia and melanoma (13,14).

Background References

- Hermann, A. et al. (2004) *Cell Mol Life Sci* 61, 2571-87.
- Turek-Plewa, J. and Jagodziński, P.P. (2005) *Cell Mol Biol Lett* 10, 631-47.
- Okano, M. et al. (1999) *Cell* 99, 247-57.
- Li, E. et al. (1992) *Cell* 69, 915-26.
- Tahiliani, M. et al. (2009) *Science* 324, 930-5.
- He, Y.F. et al. (2011) *Science* 333, 1303-7.
- Ito, S. et al. (2011) *Science* 333, 1300-3.
- Kriaucionis, S. and Heintz, N. (2009) *Science* 324, 929-30.
- Globisch, D. et al. (2010) *PLoS One* 5, e15367.
- Gao, Y. et al. (2013) *Cell Stem Cell* 12, 453-69.
- Mellén, M. et al. (2012) *Cell* 151, 1417-30.
- Wen, L. et al. (2014) *Genome Biol* 15, R49.
- Delhommeau, F. et al. (2009) *N Engl J Med* 360, 2289-301.
- Lian, C.G. et al. (2012) *Cell* 150, 1135-46.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

IF-IC: Immunofluorescence (Immunocytochemistry) **Dot Blot:** DNA Dot Blot

Cross-Reactivity Key

All: All Species Expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.