## ې Pontin/RUVBL1 (D1L8J) XP<sup>®</sup> Rabbit mAb





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, ChIP	<b>Reactivity:</b> H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 50	<b>Source/Isotype:</b> Rabbit IgG	UniProt ID: #Q9Y265	Entrez-Gene Id: 8607		
Product Usage Information	•		nal ChIP results, use 10 μl of antibody and 10 μg of chromatin (approximately 4 x 10 <sup>6</sup> cells) per ntibody has been validated using SimpleChIP <sup>®</sup> Enzymatic Chromatin IP Kits.					
		<b>Application</b> Western Blotting Chromatin IP			<b>Dilution</b> 1:1000 1:50			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less thar 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				ol and less than		
Specificity/Sensitivity		Pontin/RUVBL1 (D1L8J) XP $^{ extsf{@}}$ Rabbit mAb recognizes endogenous levels of total Pontin/RUVBL1 protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu428 of human Pontin/RUVBL1 protein.						
Background		Reptin/RuvBL2 and Pontin/RuvBL1 are closely related members of the AAA+ (ATPase associated with diverse cellular activities) superfamily of proteins, and are putatively homologous to bacterial RuvB proteins that drive branch migration of Holliday junctions (1). Reptin and Pontin function together as essential components of chromatin remodeling and modification complexes, such as INO80, TIP60, SRCAP, and Uri1, which play key roles in regulating gene transcription (1,2). In their capacity as essential transcriptional co-regulators, Reptin and Pontin have both been implicated in oncogenic transformations, including those driven by c-Myc, β-catenin, and E1A (2-7).						
Background Re	eferences	1. Jha, S. and Dutta, A. 2. Gallant, P. (2007) <i>Tr</i> . 3. Huber, O. et al. (200 4. Kim, J.H. et al. (2005 5. Bauer, A. et al. (2000 6. Wood, M.A. et al. (20 7. Dugan, K.A. et al. (2000)	<i>ends Cell Biol</i> 17, 18 8) <i>Cancer Res</i> 68, 68 6) <i>Nature</i> 434, 921-6 0) <i>EMBO J</i> 19, 6121-3 000) <i>Mol Cell</i> 5, 321	7-92. 373-6. 30. -30.				
Species Reactiv	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot B	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting ChIP: Chromatin IP						
Cross-Reactivit	ty Key	H: Human M: Mouse R: Rat Mk: Monkey						
Trademarks ar	nd Patents	Cell Signaling Technology is a trademark of Cell Signaling Tec			logy, Inc.			
		SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.						
		XP is a registered trademark of Cell Signaling Technology, Inc.						
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for		
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no						

## force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.