

**CLEC12A/CLL-1 (E8A3Z) Rabbit mAb**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H	Endogenous	55-75	Rabbit IgG	#Q5QGZ9	160364

**Product Usage Information****Application**

Western Blotting  
Immunoprecipitation

**Dilution**

1:1000  
1:100

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

CLEC12A/CLL-1 (E8A3Z) Rabbit mAb recognizes endogenous levels of total CLEC12A/CLL-1 protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human CLEC12A/CLL-1 protein.

**Background**

Part of the NK gene complex, C-type lectin-like molecule 1 (MCL/DCAL-2/CLL-1/CLEC12A) encodes a type-II transmembrane glycoprotein whose expression is largely restricted to hematopoietic cells of the myeloid lineage such as monocytes, macrophages, dendritic cells, and neutrophils (1-3). Research studies have shown that CLL-1 possesses a single C-type lectin-like domain within the extracellular domain and a single ITIM motif within its short cytoplasmic tail, which facilitates association with inhibitory SH2 domain-containing tyrosine phosphatases, SHP-1 and SHP-2. It is thought that signaling through the ITIM motif of CLL-1 facilitates inhibition of myeloid cell activation (1,2). By serving as a receptor for DAMPs that become exposed on dead cells, such as uric acid crystals, CLL-1 restrains pro-inflammatory immune responses that occur in response to cell death (4). In addition to being expressed on normal differentiated myeloid cells, research studies have also demonstrated expression of CLL-1 on the surface of malignant myeloid cells (5). As a result, CLL-1 has received significant attention as a potential novel therapeutic target for AML as its expression is absent from normal hematopoietic stem cells but is highly overexpressed on AML stem cells (5-9).

**Background References**

1. Marshall, A.S. et al. (2004) *J Biol Chem* 279, 14792-802.
2. Marshall, A.S. et al. (2006) *Eur J Immunol* 36, 2159-69.
3. Chen, C.H. et al. (2006) *Blood* 107, 1459-67.
4. Neumann, K. et al. (2014) *Immunity* 40, 389-99.
5. Bakker, A.B. et al. (2004) *Cancer Res* 64, 8443-50.
6. Jiang, Y.P. et al. (2018) *Blood Adv* 2, 1738-49.
7. Tashiro, H. et al. (2017) *Mol Ther* 25, 2202-13.
8. Wang, J. et al. (2018) *J Hematol Oncol* 11, 7.
9. Laborda, E. et al. (2017) *Int J Mol Sci* 18.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting **IP:** Immunoprecipitation

**Cross-Reactivity Key**

**H:** Human

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.