

TRPC3 (D4P5S) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	105	Rabbit IgG	#Q13507-3	7222

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TRPC3 (D4P5S) Rabbit mAb recognizes endogeneous levels of total TRPC3 protein. This antibody does not cross-react with TRPC6 or TRPC7 proteins.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human TRPC3 protein.

Background

Transient Receptor Potential Canonical 3 (TRPC3) belongs to the superfamily of TRP cation channels. The TRPC subfamily (TRPC1-7) is a group of calcium-permeable cation channels that mediates the increase in intracellular [Ca²⁺] following activation by G-protein-coupled receptors or receptor tyrosine kinases (1). TRPC3 is mainly expressed in the brain and various other tissues, though at a much lower level (2). In the brain, TRPC3 is involved in BDNF-induced axonal growth cone plasticity, dendritic spine formation, and neuronal survival (3-6). It is also required for synaptic transmission and motor coordination (7). Outside of the central nervous system, TRPC3 also exerts other important biological functions such as regulating cardiac and vascular contractility, maintaining Ca²⁺ homeostasis in primary T cells and endothelial cells (8-10). TRPC3 is activated by diacylglycerol and Inositol 1,4,5-trisphosphate (11,12). It is also activated by internal calcium store depletion and regulates mitochondrial calcium uptake (13,14).

Background References

1. Montell, C. (2005) *Sci STKE* 2005, re3.
2. Riccio, A. et al. (2002) *Brain Res Mol Brain Res* 109, 95-104.
3. Li, H.S. et al. (1999) *Neuron* 24, 261-73.
4. Li, Y. et al. (2005) *Nature* 434, 894-8.
5. Amaral, M.D. and Pozzo-Miller, L. (2007) *J Neurosci* 27, 5179-89.
6. Jia, Y. et al. (2007) *Nat Neurosci* 10, 559-67.
7. Hartmann, J. et al. (2008) *Neuron* 59, 392-8.
8. Nilius, B. et al. (2003) *Endothelium* 10, 5-15.
9. Yeon, S.I. et al. (2014) *PLoS One* 9, e110413.
10. Wenning, A.S. et al. (2011) *Biochim Biophys Acta* 1813, 412-23.
11. Hofmann, T. et al. (1999) *Nature* 397, 259-63.
12. Song, T. et al. (2015) *Am J Physiol Lung Cell Mol Physiol* 309, L1455-66.
13. Ma, H.T. et al. (2000) *Science* 287, 1647-51.
14. Feng, S. et al. (2013) *Proc Natl Acad Sci U S A* 110, 11011-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.