

SORL1 (D8D4G) Rabbit mAb



Orders: 877-616-CELL (2355) orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 250	Source/Isotype: Rabbit IgG	UniProt ID: #Q92673	Entrez-Gene Id: 6653
Product Usage Information	2	Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		SORL1 (D8D4G) Rabbit mAb recognizes endogenous levels of total SORL1 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu267 of human SORL1 protein.				
Background		SORL1, also called SorLA, is a 250 kDa type-1 membrane protein found in neurons of the central and peripheral nervous system (1). SORL1 is related to the yeast transporter Vps10p (2). While its exact function is unknown, because it shuttles between the Golgi apparatus and endolysosomal compartments, and given its structural similarity with Vps10p, it has been suggested that SORL1 acts in intracellular trafficking (3). Interestingly, SORL1 expression has been shown to be decreased in the brain of patients with Alzheimer's disease (4), and research studies have demonstrated a role for SORL1 as a neuronal sorting receptor that binds APP and regulates its trafficking and proteolytic processing, thus regulating the production of amyloid-beta peptides (5).				
Background R	eferences	1. Yamazaki, H. et al. (1996) <i>J Biol Chem</i> 271, 24761-8. 2. Marcusson, E.G. et al. (1994) <i>Cell</i> 77, 579-86. 3. Jacobsen, L. et al. (2002) <i>FEBS Lett</i> 511, 155-8. 4. Scherzer, C.R. et al. (2004) <i>Arch Neurol</i> 61, 1200-5. 5. Andersen, O.M. et al. (2005) <i>Proc Natl Acad Sci U S A</i> 102, 13461-6.				
Species Reacti	vity	Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Mantaun Diet Buffen		TADODTANT F				

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human M: Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any

commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.