

β-Actin (13E5) Rabbit mAb (Alexa Fluor® 555 Conjugate)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/26/16

For Research Use Only. Not For Use In Diagnostic Procedures.



Applications	Species Cross-Reactivity*	Isotype
IF-IC Endogenous	H, M, R, Mk, B, Pg, (C, Dg, Hr)	Rabbit IgG

Description: This Cell Signaling Technology antibody is conjugated to Alexa Fluor® 555 fluorescent dye and tested in-house for immunofluorescent analysis in monkey cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated β-Actin (13E5) Rabbit mAb #4970.

Background: Actin, a ubiquitous eukaryotic protein, is the major component of the cytoskeleton. At least six isoforms are known in mammals. Nonmuscle β- and γ-actin, also known as cytoplasmic actin, are predominantly expressed in nonmuscle cells, controlling cell structure and motility (1). α-cardiac and α-skeletal actin are expressed in striated cardiac and skeletal muscles, respectively; two smooth muscle actins, α- and γ-actin, are found primarily in vascular smooth muscle and enteric smooth muscle, respectively. These actin isoforms regulate the contractile potential of muscle cells (1). Actin exists mainly as a fibrous polymer, F-actin. In response to cytoskeletal reorganizing signals during processes such as cytokinesis, endocytosis, or stress, cofilin promotes fragmentation and depolymerization of F-actin, resulting in an increase in the monomeric globular form, G-actin (2). The Arp2/3 complex stabilizes F-actin fragments and promotes formation of new actin filaments (2). Research studies have shown that actin is hyperphosphorylated in primary breast tumors (3). Cleavage of actin under apoptotic conditions has been observed *in vitro* and in cardiac and skeletal muscle, as shown in research studies (4-6). Actin cleavage by caspase-3 may accelerate ubiquitin/proteasome-dependent muscle proteolysis (6).

Specificity/Sensitivity: β-Actin (13E5) Rabbit mAb (Alexa Fluor® 555 Conjugate) detects endogenous levels of total β-actin protein. This antibody may cross-react with the γ-actin (cytoplasmic isoform). It does not cross-react with α-skeletal, α-cardiac, α-vascular smooth, or γ-enteric smooth muscle isoforms.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino-terminus of human β-actin protein.



Confocal immunofluorescent analysis of COS-7 cells, at high confluency (upper) or low confluency (lower), using β-Actin (13E5) Rabbit mAb (Alexa Fluor® 555 Conjugate) (red). Blue pseudocolor = DRAQ5® #4084 (fluorescent DNA dye).

Entrez-Gene ID #60
UniProt ID #P60709

Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.

***Species cross-reactivity other than monkey is determined by western blot using the unconjugated antibody.**

Recommended Antibody Dilutions:
 Immunofluorescence (IF-IC) 1:50

For product specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

- Background References:**
- (1) Herman, I.M. (1993) *Curr. Opin. Cell Biol.* 5, 48-55.
 - (2) Condeelis, J. (2001) *Trends Cell Biol.* 11, 288-293.
 - (3) Lim, Y.P. et al. (2004) *Clin. Cancer Res.* 10, 3980-3987.
 - (4) Kayalar, C. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93, 2234-2238.
 - (5) Communal, C. et al. (2002) *Proc. Natl. Acad. Sci. USA.* 99, 6252-6256.
 - (6) Du, J. et al. (2004) *J. Clin. Invest.* 113, 115-123.

U.S. Patent No. 5,675,063

DRAQ5® is a registered trademark of Biostatus Limited.
 Alexa Fluor® is a registered trademark of Molecular Probes, Inc.