Store at -20C

41

ò

#

Cell Signaling UCHL3 (D25E6) Rabbit mAb H. 877-616-CELL (2355) orders@cellsignal.com Orders: Support: 877-678-TECH (8324) info@cellsignal.com cellsignal.com Web: 3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 27	Source/Isotype: Rabbit IgG	UniProt ID: #P15374	Entrez-Gene Id: 7347
Product Usage Information		Application Western Blotting		Dilution 1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		UCHL3 (D25E6) Rabbit mAb recognizes endogenous levels of total UCHL3 protein. This antibody does not cross-react with either UCHL1 or BAP1 and is not predicted to cross-react with UCHL5.				
Species predicted to react based on 100% sequence homology		Bovine, Dog, Horse				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human UCHL3 protein.				
Background		Protein ubiquitination and deubiquitination are reversible processes catalyzed by ubiquitinating enzymes (UBEs) and deubiquitinating enzymes (DUBs) (1,2). DUBs are categorized into 5 subfamilies: USP, UCH, OTU, MJD, and JAMM. UCHL1, UCHL3, UCHL5/UCH37, and BRCA-1-associated protein-1 (BAP1) belong to the UCH family of DUBs, which all posses a conserved catalytic domain (UCH domain) of about 230 amino acids. UCHL5 and BAP1 have unique extended C-terminal tails. UCHL1 is abundantly expressed in neuronal tissues and testes, while UCHL3 expression is more widely distributed (3,4). Although UCHL1 and UCHL3 are the most closely related UCH family members with about 53% identity, their biochemical properties differ in that UCHL1 binds monoubiquitin and UCHL3 shows dual specificity toward both ubiquitin (Ub) and NEDD8, a Ub-like molecule. In particular, UCHL3 functions as a Ub hydrolase involved in the processing of both Ub precursors and ubiquitinated substrates, generating free monomeric Ub. This is accomplished through the ability of UCHL3 to recognize and hydrolyze isopeptide bonds at the C-terminal glycine of either Ub or NEDD8 (5-7). Recent functional studies have identified UCH-L3 as a critical regulator of adipogenesis through its ability to promote IGF-IR and insulin receptor signaling (8). Furthermore, UCHL3 has been shown to promote deubiquitination, recycling, and cell surface expression of the epithelial sodium channel (9).				
Background Re	ferences	1. Nijman, S.M. et al. (2 2. Nalepa, G. et al. (20 3. Leroy, E. et al. (1998 4. Kurihara, L.J. et al. (20 5. Osaka, H. et al. (200 6. Wada, H. et al. (1999 7. Kwon, J. (2007) <i>Exp</i> 8. Suzuki, M. et al. (200 9. Butterworth, M.B. e	06) <i>Nat Rev Drug D 1) Nature</i> 395, 451-2 2001) <i>Hum Mol Gei</i> 13) <i>Hum Mol Genet 8) Biochem Biophy</i> <i>Anim</i> 56, 71-7. 09) <i>Endocrinology</i> 1	<i>iscov</i> 5, 596-613. 2. net 10, 1963-70. 12, 1945-58. 5 <i>Res Commun</i> 251, 688- 150, 5230-9.	92.	
Species Reactiv	vity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.