ò

Acetyl-α-Tubulin (Lys40) (D20G3) XP[®] Rabbit mAb (Alexa Fluor[®] 647 Conjugate)



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IF-IC, FC-FP	Reactivity: H M R Mk Z	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P68363	Entrez-Gene Id: 10376	
Product Usage Information		Application Immunofluorescence (Immunocytochemistry) Flow Cytometry (Fixed/Permeabilized)		Dilution 1:50 1:50		
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.				
Specificity/Sensit	tivity	Acetyl- α -Tubulin (Lys40) (D20G3) XP [®] Rabbit mAb (Alexa Fluor [®] 647 Conjugate) detects endogenous levels of α -tubulin only when acetylated at Lys40. This amino acid is not conserved in β -tubulin.				
Species predicted based on 100% se homology	d to react equence	Xenopus				
Source / Purificat	tion	Monoclonal antibody is produced by immunizing animals with a synthetic acetylpeptide corresponding to residues surrounding Lys40 of human α-tubulin.				
Description		This Cell Signaling Technology antibody is conjugated to Alexa Fluor [®] 647 fluorescent dye and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Acetyl-α-Tubulin (Lys40) (D20G3) XP [®] Rabbit mAb #5335.				
Background		The cytoskeleton consists of three types of cytosolic fibers: microtubules, microfilaments (actin filaments), and intermediate filaments. Globular tubulin subunits comprise the microtubule building block, with α/β -tubulin heterodimers forming the tubulin subunit common to all eukaryotic cells. γ -tubulin is required to nucleate polymerization of tubulin subunits to form microtubule polymers. Many cell movements are mediated by microtubule action, including the beating of cilia and flagella, cytoplasmic transport of membrane vesicles, chromosome alignment during meiosis/mitosis, and nerve-cell axon migration. These movements result from competitive microtubule polymerization and depolymerization or through the actions of microtubule motor proteins (1).				
Background Refe	erences	1. Westermann, S. and Weber, K. (2003) Nat Rev Mol Cell Biol 4, 938-47.				
Species Reactivit	-	Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Applications Key		IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized)				
Cross-Reactivity	Кеу	H: Human M: Mouse R: Rat Mk: Monkey Z: Zebrafish				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		_	d trademark of Life Techn			
		transfer of this product is conducted by the buyer, must not (1) use this pro- purposes; (b) testing, and per-test basis; or (c) man this product or its compo- purchasing a license to th	s conditioned on the buye excluding contract resear duct or its components fo alysis or screening service ufacturing or quality assu- nents for resale, whether his product for purposes	er using the purchase ch or any fee for serv r (a) diagnostic, ther es, or information in arance or quality con or not resold for us other than as describ	return for compensation on a trol, and/or (2) sell or transfer e in research. For information on	

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.