Store at +4C

8183

#

Phospho-Stat1 (Tyr701) (58D6) Rabbit mAb (Alexa Fluor[®] 555 Conjugate)



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IF-IC, FC-FP	Reactivity: H M	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P42224	Entrez-Gene Id: 6772		
Product Usage Information		Application Immunofluorescence (Immunocytochemistry) Flow Cytometry (Fixed/Permeabilized)			Dilution 1:50 1:50		
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.					
Specificity/Sensit	tivity	Phospho-Stat1 (Tyr701) (58D6) Rabbit mAb (Alexa Fluor [®] 555 Conjugate) detects endogenous levels of Stat1 only when phosphorylated at Tyr701. The antibody detects phosphorylated Tyr701 of both p91 and p84 Stat1. It does not cross-react with the corresponding phospho-tyrosines of other Stat proteins.					
Source / Purificat	tion	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr701 of human Stat1.					
Description		This Cell Signaling Technology antibody is conjugated to Alexa Fluor [®] 555 fluorescent dye and tested in-house for immunofluroescent analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-Stat1 (Tyr701) (58D6) Rabbit mAb #9167.					
Background		The Stat1 transcription factor is activated in response to a large number of ligands (1) and is essential for responsiveness to IFN- α and IFN- γ (2,3). Phosphorylation of Stat1 at Tyr701 induces Stat1 dimerization, nuclear translocation, and DNA binding (4). Stat1 protein exists as a pair of isoforms, Stat1 α (91 kDa) and the splice variant Stat1 β (84 kDa). In most cells, both isoforms are activated by IFN- α , but only Stat1 α is activated by IFN- γ . The inappropriate activation of Stat1 occurs in many tumors (5). In addition to tyrosine phosphorylation, Stat1 is also phosphorylated at Ser727 through a p38 mitogenactivated protein kinase (MAPK)-dependent pathway in response to IFN- α and other cellular stresses (6). Serine phosphorylation may be required for the maximal induction of Stat1-mediated gene activation.					
Background Refe	erences	1. Heim, M.H. (1999) <i>J Recept Signal Transduct Res</i> 19, 75-120. 2. Durbin, J.E. et al. (1996) <i>Cell</i> 84, 443-50. 3. Meraz, M.A. et al. (1996) <i>Cell</i> 84, 431-42. 4. Ihle, J.N. et al. (1994) <i>Trends Biochem Sci</i> 19, 222-7. 5. Frank, D.A. (1999) <i>Mol Med</i> 5, 432-56. 6. Wen, Z. et al. (1995) <i>Cell</i> 82, 241-50.					
Species Reactivit	у	Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Applications Key		IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized)					
Cross-Reactivity	Key	H: Human M: Mouse					
Trademarks and	Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.					
		transfer of this product is conducted by the buyer, must not (1) use this pro purposes; (b) testing, and per-test basis; or (c) man this product or its compo purchasing a license to th	s conditioned on the buye excluding contract resear duct or its components fo alysis or screening service ufacturing or quality assu- nents for resale, whether his product for purposes	er using the purchase ch or any fee for serv or (a) diagnostic, thera es, or information in r urance or quality cont or not resold for use other than as describ	eturn for compensation on a trol, and/or (2) sell or transfer e in research. For information on		

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.