

Store at  
-20°C

#84200

# Steroid Hormone Receptor Antibody Sampler Kit



**Support:** +1-978-867-2388 (U.S.)  
www.cellsignal.com/support

**Orders:** 877-616-2355 (U.S.)  
orders@cellsignal.com

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Product Includes	Product #	Quantity	Mol. Wt.	Isotype/Source
Androgen Receptor (D6F11) XP <sup>®</sup> Rabbit mAb	5153	20 µL	110 kDa	Rabbit IgG
Estrogen Receptor α (D8H8) Rabbit mAb	8644	20 µL	66 kDa	Rabbit IgG
Glucocorticoid Receptor (D6H2L) XP <sup>®</sup> Rabbit mAb	12041	20 µL	94, 91 kDa	Rabbit IgG
Progesterone Receptor A/B (D8Q2J) XP <sup>®</sup> Rabbit mAb	8757	20 µL	90 (PR-A), 118 (PR-B) kDa	Rabbit IgG
Mineralocorticoid Receptor (E9W1M) Rabbit mAb	58883	20 µL	120 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µL		Goat

See [www.cellsignal.com](http://www.cellsignal.com) for individual component applications, species cross-reactivity, dilutions, and additional application protocols.

**Description:** The Steroid Hormone Receptor Antibody Sampler Kit provides an economical means of detecting levels of steroid hormone nuclear receptors. The kit includes enough antibodies to perform two western blot experiments with each primary antibody.

**Background:** Androgen receptor (AR), a zinc finger transcription factor belonging to the nuclear receptor superfamily, is activated by phosphorylation and dimerization upon ligand binding (1). This promotes nuclear localization and binding of AR to androgen response elements in androgen target genes. Research studies have shown that AR plays a crucial role in several stages of male development and the progression of prostate cancer (2,3). Estrogen receptor α (ERα), a member of the steroid receptor superfamily, contains highly conserved DNA-binding and ligand-binding domains (4). Through its estrogen-independent and estrogen-dependent activation domains (AF-1 and AF-2, respectively), ERα regulates transcription by recruiting coactivator proteins and interacting with general transcriptional machinery (5). Human progesterone receptor (PR) is expressed as two forms: the full length PR-B and the short form PR-A. PR-A lacks the first 164 amino acid residues of PR-B (6,7). Both PR-A and PR-B are ligand activated, but differ in their relative ability to activate target gene transcription (8,9). Glucocorticoid hormones control cellular proliferation, inflammation, and metabolism through their association with the glucocorticoid receptor (GR)/NR3C1, a member of the nuclear hormone receptor superfamily of transcription factors (10). GR is composed of several conserved structural elements, including a carboxy-terminal ligand-binding domain (which also contains residues critical for receptor dimerization and hormone-dependent gene transactivation), a neighboring hinge region containing nuclear localization signals, a central zinc-finger-containing DNA-binding domain, and an amino-terminal variable region that participates in ligand-independent gene transcription. In the absence of hormone, a significant population of GR is localized to the cytoplasm in an inactive form via its association with regulatory chaperone proteins, such as HSP90, HSP70, and FKBP52. On hormone binding,

GR is released from the chaperone complex and translocates to the nucleus as a dimer to associate with specific DNA sequences termed glucocorticoid response elements (GREs), thereby enhancing or repressing transcription of specific target genes (11). Mineralocorticoid receptor (MR) is a steroid hormone receptor with structural and functional similarities to GR. MR binds with high affinity to aldosterone and other mineralocorticoids as well as glucocorticoids (12-14). Upon ligand binding, MR undergoes conformational changes and enters the nucleus to bind to target mineralocorticoid response elements (MREs) (4,15,16). MR is also able to heterodimerize with GR and bind to hormone response elements on DNA in cells that express both receptors (17-19).

**Specificity/Sensitivity:** Each antibody in the Steroid Hormone Receptor Antibody Sampler Kit detects endogenous levels of its target protein. Androgen Receptor (D6F11) XP<sup>®</sup> Rabbit mAb detects endogenous levels of total androgen receptor protein. Estrogen Receptor α (D8H8) Rabbit mAb recognizes endogenous levels of total ERα protein. Progesterone Receptor A/B (D8Q2J) XP<sup>®</sup> Rabbit mAb recognizes endogenous levels of total progesterone receptor A and B proteins. This antibody does not cross-react with either the glucocorticoid receptor or the mineralocorticoid receptor. Glucocorticoid Receptor (D6H2L) XP<sup>®</sup> Rabbit mAb recognizes endogenous levels of total GR protein. This antibody reacts with GR-α and GR-β but does not cross-react with mineralocorticoid receptor. Mineralocorticoid Receptor (E9W1M) Rabbit mAb recognizes endogenous levels of total mineralocorticoid receptor protein.

**Source/Purification:** Monoclonal antibodies are produced by immunizing animals with recombinant proteins corresponding to residues near the amino terminal region of human androgen receptor protein, specific to the amino terminus of human GR protein and human mineralocorticoid receptor protein; with synthetic peptides corresponding to residues in the carboxy terminus of human ERα protein and surrounding Tyr541 of human progesterone receptor protein.

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/mL BSA, 50% glycerol, and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibodies.

Please visit [www.cellsignal.com](http://www.cellsignal.com) for validation data and a complete listing of recommended companion products.

#### Background References:

- (1) Li, J. and Al-Azzawi, F. (2009) *Maturitas* 63, 142-8.
- (2) Avila, D.M. et al. (2001) *J Steroid Biochem Mol Biol* 76, 135-42.
- (3) Montgomery, J.S. et al. (2001) *J Pathol* 195, 138-46.
- (4) Mangelsdorf, D.J. et al. (1995) *Cell* 83, 835-9.
- (5) Glass, C.K. and Rosenfeld, M.G. (2000) *Genes Dev* 14, 121-41.
- (6) Evans, R.M. (1988) *Science* 240, 889-95.
- (7) Kastner, P. et al. (1990) *EMBO J* 9, 1603-14.
- (8) Giangrande, P.H. et al. (2000) *Mol Cell Biol* 20, 3102-15.
- (9) Wen, D.X. et al. (1994) *Mol Cell Biol* 14, 8356-64.
- (10) Yamamoto, K.R. (1985) *Annu Rev Genet* 19, 209-52.
- (11) Necela, B.M. and Cidlowski, J.A. (2003) *Trends Pharmacol Sci* 24, 58-61.
- (12) Arriza, J.L. et al. (1987) *Science* 237, 268-75.
- (13) Giguère, V. et al. (1988) *Nature* 331, 91-4.
- (14) Beato, M. et al. (1995) *Cell* 83, 851-7.
- (15) Guiochon-Mantel, A. et al. (1996) *J Steroid Biochem Mol Biol* 56, 3-9.
- (16) Liu, W. et al. (1995) *Proc Natl Acad Sci U S A* 92, 12480-4.
- (17) Liu, W. et al. (1996) *Mol Endocrinol* 10, 1399-406.
- (18) Trapp, T. et al. (1994) *Neuron* 13, 1457-62.
- (19) Funder, J.W. (1995) *J Steroid Biochem Mol Biol* 53, 53-5.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

Thank you for your recent purchase. If you would like to provide a review visit [cellsignal.com/comments](http://cellsignal.com/comments).

[www.cellsignal.com](http://www.cellsignal.com)

© 2022 Cell Signaling Technology, Inc.

XP and Cell Signaling Technology are trademarks of Cell Signaling Technology, Inc.

**Applications:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide **Species Cross-Reactivity:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.