

#8487
 Store at -20C

Sestrin-2 (D1B6) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H Mk	Endogenous	56	Rabbit IgG	#P58004	83667

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Sestrin-2 (D1B6) Rabbit mAb recognizes overexpressed levels of total sestrin-2 protein, as well as endogenous levels of sestrin-2 protein when expression is high.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human sestrin-2 protein.

Background

The Sestrin family of proteins, which comprises sestrin-1 (PA26), sestrin-2 (Hi95), and sestrin-3, plays an important role in regulating the cellular response to oxidative stress. Sestrin-1 and -2, both transcriptional targets of p53, have been shown to activate AMPK activity toward TSC2, ultimately resulting in the inhibition of mTOR (1). Thus, sestrin-1 and -2 link p53 signaling to the regulation of cellular growth (2). Sestrins have also been shown to function as modulators of cellular hydrogen peroxide concentration. Hydrogen peroxide is an important signaling molecule that is highly genotoxic and must therefore be tightly regulated (3). 2-Cys peroxiredoxins are metabolizing enzymes that maintain low cellular concentrations of hydrogen peroxide, but are hyperoxidized to an inactive state by a rapid increase in concentration (4) before quickly being reduced back to their active state by sestrins (5). This allows a hydrogen peroxide signal to temporarily circumvent the protective activity of 2-Cys peroxiredoxins, while minimizing oxidative damage.

Background References

1. Budanov, A.V. and Karin, M. (2008) *Cell* 134, 451-60.
2. Wullschleger, S. et al. (2006) *Cell* 124, 471-84.
3. Veal, E.A. et al. (2007) *Mol Cell* 26, 1-14.
4. Jönsson, T.J. and Lowther, W.T. (2007) *Subcell Biochem* 44, 115-41.
5. Budanov, A.V. et al. (2004) *Science* 304, 596-600.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.