



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at +4C
#8520

Phospho-S6 Ribosomal Protein (Ser235/236) (D57.2.2E) XP[®] Rabbit mAb (Pacific Blue[™] Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M R Mk Mi Sc	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P62753	Entrez-Gene Id: 6194
-------------------------------	--------------------------------------	-----------------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Flow Cytometry (Fixed/Permeabilized)	Dilution 1:50
Storage	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.	
Specificity/Sensitivity	Phospho-S6 Ribosomal Protein (Ser235/236) (D57.2.2E) XP [®] Rabbit mAb (Pacific Blue [™] Conjugate) recognizes endogenous levels of ribosomal protein S6 only when phosphorylated at Ser235 and Ser236.	
Species predicted to react based on 100% sequence homology	Chicken, Pig	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser235 and Ser236 of human ribosomal protein S6.	
Description	This Cell Signaling Technology antibody is conjugated to Pacific Blue [™] fluorescent dye and tested in-house for direct flow cytometry in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-S6 Ribosomal Protein (Ser235/236) (D57.2.2E) XP [®] Rabbit mAb #4858.	
Background	One way that growth factors and mitogens effectively promote sustained cell growth and proliferation is by upregulating mRNA translation (1,2). Growth factors and mitogens induce the activation of p70 S6 kinase and the subsequent phosphorylation of S6 ribosomal protein. Phosphorylation of S6 ribosomal protein correlates with an increase in translation of mRNA transcripts that contain an oligopyrimidine tract in their 5' untranslated regions (2). These particular mRNA transcripts (5'TOP) encode proteins involved in cell cycle progression, as well as ribosomal proteins and elongation factors necessary for translation (2,3). Important S6 ribosomal protein phosphorylation sites include several residues (Ser235, Ser236, Ser240, and Ser244) located within a small, carboxy-terminal region of S6 protein (4,5).	
Background References	<ol style="list-style-type: none"> 1. Dufner, A. and Thomas, G. (1999) <i>Exp Cell Res</i> 253, 100-9. 2. Peterson, R.T. and Schreiber, S.L. (1998) <i>Curr Biol</i> 8, R248-50. 3. Jefferies, H.B. et al. (1997) <i>EMBO J</i> 16, 3693-704. 4. Ferrari, S. et al. (1991) <i>J Biol Chem</i> 266, 22770-5. 5. Flotow, H. and Thomas, G. (1992) <i>J Biol Chem</i> 267, 3074-8. 	

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key **H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey **Mi:** Mink **Sc:** S. cerevisiae

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Pacific Blue is a trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.