

NKCC1 (D2O8R) XP[®] Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IF-F	Reactivity: M R	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P55012	Entrez-Gene Id: 20496
------------------------------	---------------------------	-----------------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information**Application**

Immunofluorescence (Frozen)

Dilution

1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

NKCC1 (D2O8R) XP[®] Rabbit mAb recognizes endogenous levels of total NKCC1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of mouse NKCC1 protein.

Background

The electroneutral cation-chloride-coupled co-transporter (SLC12) gene family comprises bumetanide-sensitive Na⁺/K⁺/Cl⁻ (NKCC), thiazide-sensitive Na⁺/Cl⁻, and K⁺/Cl⁻ (KCC) co-transporters. SLC12A1/NKCC2 and SLC12A2/NKCC1 regulate cell volume and maintain cellular homeostasis in response to osmotic and oxidative stress (1). The broadly expressed NKCC1 is thought to play roles in fluid secretion (i.e. salivary gland function), salt balance (i.e. maintenance of renin and aldosterone levels), and neuronal development and signaling (2-7). During neuronal development, NKCC1 and KCC2 maintain a fine balance between chloride influx (NKCC1) and efflux (KCC2), which regulates γ-aminobutyric acid (GABA)-mediated neurotransmission (3). Increased NKCC1 expression in immature neurons maintains high intracellular chloride levels that result in inhibitory GABAergic signaling; KCC2 maintains low intracellular chloride levels and excitatory GABAergic responses in mature neurons (4,5,8). Deletion of NKCC1 impairs NGF-mediated neurite outgrowth in PC-12D cells while inhibition of NKCC1 with bumetanide inhibits re-growth of axotomized dorsal root ganglion cells (6,7). Defective chloride homeostasis in neurons is linked to seizure disorders that are ameliorated by bumetanide treatment, indicating that abnormal NKCC1 and NKCC2 expression or signaling may play a role in neonatal and adult seizures (9-12). NKCC1 is found as a homodimer or within heterooligomers with other SLC12 family members. This transport protein associates with a number of oxidative- and osmotic-responsive kinases that bind, phosphorylate, and activate NKCC1 co-transporter activity (13-16). In response to decreased intracellular chloride concentrations, Ste20-related proline-alanine-rich kinase (SPAK) phosphorylates NKCC1 to increase co-transporter activity and promote chloride influx (16-19). Oxidative stress response kinase 1 (OSR1) also phosphorylates and activates NKCC1 in response to oxidative stress (14).

Background References

1. Hebert, S.C. et al. (2004) *Pflugers Arch* 447, 580-93.
2. Evans, R.L. et al. (2000) *J Biol Chem* 275, 26720-6.
3. Kim, S.M. et al. (2008) *Am J Physiol Renal Physiol* 295, F1230-8.
4. Khirug, S. et al. (2008) *J Neurosci* 28, 4635-9.
5. Kahle, K.T. et al. (2008) *Nat Clin Pract Neurol* 4, 490-503.
6. Nakajima, K. et al. (2007) *Biochem Biophys Res Commun* 359, 604-10.
7. Pieraut, S. et al. (2007) *J Neurosci* 27, 6751-9.
8. Ben-Ari, Y. (2002) *Nat Rev Neurosci* 3, 728-39.
9. Fukuda, A. (2005) *Nat Med* 11, 1153-4.
10. Dzhalal, V.I. et al. (2005) *Nat Med* 11, 1205-13.
11. Jayakumar, A.R. et al. (2008) *J Biol Chem* 283, 33874-82.
12. Kahle, K.T. and Staley, K.J. (2008) *Neurosurg Focus* 25, E22.
13. Moore-Hoon, M.L. and Turner, R.J. (2000) *Biochemistry* 39, 3718-24.
14. Simard, C.F. et al. (2007) *J Biol Chem* 282, 18083-93.
15. Piechotta, K. et al. (2002) *J Biol Chem* 277, 50812-9.
16. Dowd, B.F. and Forbush, B. (2003) *J Biol Chem* 278, 27347-53.
17. Geng, Y. et al. (2009) *J Biol Chem* 284, 14020-8.
18. Smith, L. et al. (2008) *J Biol Chem* 283, 22147-56.
19. Gagnon, K.B. et al. (2006) *Mol Cell Biol* 26, 689-98.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

IF-F: Immunofluorescence (Frozen)

Cross-Reactivity Key

M: Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.