



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#8611

## TIAR (D26E4) Rabbit mAb

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IP, eCLIP	<b>Reactivity:</b> H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 42	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q01085	<b>Entrez-Gene Id:</b> 7073
--------------------------------------	--------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	--------------------------------

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation  
eCLIP

#### Dilution

1:1000  
1:100  
1:200

For more information about the RBP-eCLIP service please visit Eclipsebio.

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

TIAR (D26E4) Rabbit mAb recognizes endogenous levels of total TIAR protein. This antibody also weakly recognizes recombinant TIA1 protein.

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro337 of human TIAR protein.

### Background

TIAR is a member of the RNA-recognition motif (RRM) family of RNA-binding proteins (1,2). It functions as a translational repressor under conditions of cellular damage (3,4). In response to cellular stress, TIAR associates with eIF1, eIF3, and the 40S ribosomal subunit and forms noncanonical preinitiation complexes that are translationally inactive (3,4). TIAR then aggregates with its family member TIA1 and facilitates the accumulation of the translationally inactive preinitiation complexes into discrete cytoplasmic foci called stress granules. The two major isoforms of TIAR are the products of alternative mRNA splicing (5,6).

### Background References

1. Tian, Q. et al. (1991) *Cell* 67, 629-39.
2. Kawakami, A. et al. (1992) *Proc Natl Acad Sci USA* 89, 8681-5.
3. Anderson, P. and Kedersha, N. (2002) *Cell Stress Chaperones* 7, 213-21.
4. Anderson, P. and Kedersha, N. (2002) *J Cell Sci* 115, 3227-34.
5. Kawakami, A. et al. (1994) *J Immunol* 152, 4937-45.
6. Beck, A.R. et al. (1996) *Nucleic Acids Res* 24, 3829-35.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IP:** Immunoprecipitation **eCLIP:** eCLIP

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.