

#86628 Store at +4C

MHC Class II (I-A/I-E) (M5/114.15.2) Rat mAb (violetFluor 450 Conjugate)



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
FC-FP, FC-L	M	Endogenous	Rat IgG2b kappa	#P18468, #P06342, #Q3U060, #P14435	14969, 14961, 381091, 14960

Product Usage Information

For optimal flow cytometry results, we recommend 0.25 µg of antibody per test.

Application	Dilution
Flow Cytometry (Fixed/Permeabilized)	1:80
Flow Cytometry (Live)	1:80

Storage

Supplied in 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% NaN₃, 0.1% gelatin, pH 7.2. This product is stable for 12 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.

Specificity/Sensitivity

MHC Class II (I-A/I-E) (M5/114.15.2) Rat mAb (violetFluor 450 Conjugate) recognizes endogenous levels of total MHC class II (I-A/I-E) proteins. This antibody detects epitopes within the extracellular domain of MHC class II (I-A/I-E).

Source / Purification

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.

Description

This Cell Signaling Technology antibody is conjugated to violetFluor 450 and tested in-house for direct flow cytometric analysis in mouse cells.

Background

Major histocompatibility complex class II (MHC class II) molecules are heterodimeric, transmembrane glycoproteins expressed on the surface of antigen-presenting cells, such as macrophages, dendritic cells, and B cells. Expression can also be induced on other cell types through interferon-γ signaling (1). Prior to being displayed on the cell membrane, MHC class II molecules are loaded with exogenous peptide antigens approximately 15-24 amino acids in length that were derived from endocytosed extracellular proteins digested in the lysosome (2). Antigen-presentation through MHC class II is required for T cell activation during the immune response to extracellular pathogens (2). In humans, the MHC class II protein complex is encoded by the human leukocyte antigen gene complex (HLA). HLAs corresponding to MHC class II are HLA-DP, HLA-DM, HLA-DOA, HLA-DOB, HLA-DQ, and HLA-DR (3).

The M5/114.15.2 antibody reacts with mouse MHC class II, both I-A and I-E subregion-encoded glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). It detects a polymorphic determinant present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes, but not from mice carrying the H-2s or H-2f haplotypes (4-7). The M5/114 mAb is reported to inhibit I-A-restricted T cell responses of the H-2b, H-2d, H-2q, and H-2u haplotypes, but not H-2f, H-2k, or H-2s haplotypes (8,9).

Background References

1. Ting, J.P. and Trowsdale, J. (2002) *Cell* 109 Suppl, S21-33.
2. Cresswell, P. (1994) *Annu Rev Immunol* 12, 259-93.
3. Karp, D.R. et al. (1990) *J Exp Med* 171, 615-28.
4. Staehli, F. et al. (2012) *J Immunol* 188, 3820-8.
5. Scarlett, U.K. et al. (2012) *J Exp Med* 209, 495-506.
6. Anderson, M.S. and Miller, J. (1992) *Proc Natl Acad Sci U S A* 89, 2282-6.
7. Miyazaki, T. et al. (1996) *Cell* 84, 531-41.
8. Parra, D. et al. (2012) *J Leukoc Biol* 91, 525-36.
9. Chen, M. et al. (2011) *J Immunol* 187, 5684-92.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

FC-FP: Flow Cytometry (Fixed/Permeabilized) **FC-L:** Flow Cytometry (Live)

Cross-Reactivity Key

M: Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.