

HO-1 (E9H3A) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

info@cellsignal.com cellsignal.com Web:

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-F, IF-IC	Reactivity: M	Sensitivity: Endogenous	MW (kDa): 28	Source/Isotype: Rabbit IgG	UniProt ID: #P14901	Entrez-Gene Id: 15368
Product Usage Information	•	Application Western Blotting Immunoprecipitation Immunofluorescence Immunofluorescence	(Frozen)	istry)		Dilution 1:1000 1:50 1:1600 1:1600
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		HO-1 (E9H3A) Rabbit mAb recognizes endogenous levels of total HO-1 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu251 of mouse HO-1 protein.				
Background		Heme oxygenase (HO) is the rate-limiting enzyme in the catabolism of heme that results in the release of carbon monoxide, iron, and biliverdin (1). The products of this enzymatic reaction play important biological roles in antioxidant, anti-inflammatory, and cytoprotective functions (2). HO comprises two isozymes, including the constitutively expressed HO-2 isozyme and the inducible HO-1 isozyme (3). Inducible HO-1 is expressed as an adaptive response to several stimuli, including heme, metals, and hormones (4). The induction of HO-1 has been implicated in numerous disease states, such as transplant rejection, hypertension, atherosclerosis, Alzheimer's disease, endotoxic shock, diabetes, inflammation, and neurological disorders (1,5).				
Background References		 Abraham, N.G. and Kappas, A. (2008) <i>Pharmacol Rev</i> 60, 79-127. Otterbein, L.E. et al. (2003) <i>Trends Immunol</i> 24, 449-55. Cruse, I. and Maines, M.D. (1988) <i>J Biol Chem</i> 263, 3348-53. Maines, M.D. (1988) <i>FASEB J</i> 2, 2557-68. Schipper, H.M. et al. (2009) <i>Curr Alzheimer Res</i> 6, 424-30. 				
Species Reacti	vity	Species reactivity is d	etermined by testin	g in at least one approv	ed application (e.g.,	, western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting IP: Immunoprecipitation IF-F: Immunofluorescence (Frozen) IF-IC: Immunofluorescence (Immunocytochemistry)				
Cross-Reactivity Key		M: Mouse				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		XP is a registered trademark of Cell Signaling Technology, Inc.				
		All other trademarks more information.	are the property of	their respective owners.	Visit cellsignal.com	n/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of negative or effect.				

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.