## EphA3/A4/A5 (D2C11) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IP	Reactivity: H M	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 135	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P54756, #P29320, #P54764	<b>Entrez-Gene Id:</b> 2044, 2042, 2043
Product Usage Information		Application Western Blotting Immunoprecipitation			<b>Dilution</b> 1:1000 1:100	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		EphA3/A4/A5 (D2C11) Rabbit mAb recognizes endogenous levels of total EphA3, EphA4, and EphA5 proteins.				
Species predic based on 100% homology		Rat				
Source / Purifi	cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Tyr779 of human EphA3 protein.				
Background		The Eph receptors are the largest known family of receptor tyrosine kinases (RTKs). They can be divided				

The Eph receptors are the largest known family of receptor tyrosine kinases (RTKs). They can be divided into two groups based on sequence similarity and on their preference for a subset of ligands. While EphA receptors bind to a glycosylphosphatidylinositol-anchored ephrin A ligand, EphB receptors bind to ephrin B proteins that have a transmembrane and cytoplasmic domain (1,2). Research studies have shown that Eph receptors and ligands may be involved in many diseases including cancer (3). Both ephrin A and B ligands have dual functions. As RTK ligands, ephrins stimulate the kinase activity of Eph receptors and activate signaling pathways in receptor-expressing cells. The ephrin extracellular domain is sufficient for this function as long as it is clustered (4). The second function of ephrins has been described as "reverse signaling", whereby the cytoplasmic domain becomes tyrosine phosphorylated, allowing interactions with other proteins that may activate signaling pathways in the ligand-expressing cells (5).

The EphA3 receptor preferentially binds ephrin-A5. This ligand-receptor interaction stimulates EphA3 signaling, regulates cell adhesion and migration, and induces cellular morphologic responses (6-8). EphA3 plays a critical role in callosal axon guidance (9), retinotectal mapping of neurons (10), as well as cardiac cell migration and differentiation (11). Investigators have shown that somatic mutations in functional domains of EphA3 are linked to lung cancer progression (12). In addition, EphA3 expression levels have been correlated with tumor angiogenesis and progression in gastric and colorectal carcinoma (13,14).

## **Background References**

- 1. Wilkinson, D.G. (2000) Int Rev Cytol 196, 177-244.
- 2. Klein, R. (2001) Curr Opin Cell Biol 13, 196-203.
- 3. Dodelet, V.C. and Pasquale, E.B. (2000) *Oncogene* 19, 5614-9.
- 4. Holder, N. and Klein, R. (1999) Development 126, 2033-44.
- 5. Brückner, K. et al. (1997) *Science* 275, 1640-3.
- 6. Smith, L.M. et al. (2004) Exp Cell Res 292, 295-303.
- 7. Clifford, N. et al. (2008) J Cell Biochem 105, 1250-9.
- 8. Vearing, C. et al. (2005) *Cancer Res* 65, 6745-54.
- 9. Nishikimi, M. et al. (2011) J Neurosci 31, 16251-60.
- 10. Connor, R.J. et al. (1998) *Dev Biol* 193, 21-35.
- 11. Li, Y.Y. et al. (1998) Am J Physiol 274, H331-41.
- 12. Lisabeth, E.M. et al. (2012) Biochemistry 51, 1464-75.
- 13. Xi, H.Q. et al. (2012) *J Gastroenterol* 47, 785-94.
- 14. Xi, H.Q. and Zhao, P. (2011) *J Clin Pathol* 64, 498-503.

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat

dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

W: Western Blotting IP: Immunoprecipitation **Applications Key** 

**Cross-Reactivity Key** H: Human M: Mouse

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

**Limited Uses** Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.