

Store at
-20°C

Mouse IL-1 α Recombinant Protein




#88205


20 μ g**Support:** +1-978-867-2388 (U.S.)
www.cellsignal.com/support**Orders:** 877-616-2355 (U.S.)
orders@cellsignal.com**Entrez-Gene ID** #16175
UniProt ID #P01582

For Research Use Only. Not For Use In Diagnostic Procedures.

Background: Interleukin 1 alpha (IL-1 α) belongs to the IL-1 family of cytokines with 11 members including IL-1 β . IL-1 α is expressed in many cell types of both hematopoietic and non-hematopoietic origins under steady state, and its expression can be increased in response to appropriate stimuli (1,2). Like IL-1 β , IL-1 α is also synthesized as a precursor (pro-IL-1 α) and can be cleaved into smaller mature forms. However, both pro-IL-1 α and the cleaved form of IL-1 α are biologically active and can activate the signaling pathway through the membrane receptor IL-1R1. IL-1 α is active both as a secreted form and as a membrane-bound form. Due to such characteristics, passive leakage of IL-1 α from dying cells can activate inflammation, leading some researchers to consider IL-1 α as a key "alarmin in the cell" that alerts the host to damage or injury (3,4). In addition, IL-1 α can also enter the nucleus to modulate transcription (5,6).

Molecular Weight: 18.1 kDa**Endotoxin:** Endotoxin levels are \leq 1 EU / 1 μ g mL-1 α .**Purity:** \geq 95% purity was determined by SDS-PAGE.**Source/Purification:** Recombinant mouse IL-1 α was expressed in *E. coli* and is supplied in a lyophilized form.**Bioactivity:** The bioactivity of recombinant mL-1 α was determined in a D10.G4.1 cell proliferation assay. The ED₅₀ of each lot is \leq 10 pg/ml.

The purity of Mouse IL-1 α Recombinant Protein was determined by SDS-PAGE of 1 μ g reduced (+) and non-reduced (-) recombinant mL-1 α and staining with Coomassie Blue.



Serial dilutions of Mouse IL-1 α Recombinant Protein were added to D10.G4.1 cells. Cell proliferation was measured and the linear portion of the curve was used to calculate the ED₅₀.

Storage: Mouse IL-1 α Recombinant Protein is supplied as lyophilized material that is very stable at -20°C. It is recommended to reconstitute with sterile water at a concentration of 0.1 mg/ml which can be further diluted in aqueous solutions as needed. Addition of a carrier protein (0.1% HSA or BSA) is recommended for long-term storage.

Background References:

- (1) Garlanda, C. et al. (2013) *Immunity* 39, 1003-18.
- (2) Palomo, J. et al. (2015) *Cytokine* 76, 25-37.
- (3) Bertheloot, D. and Latz, E. (2017) *Cell Mol Immunol* 14, 43-64.
- (4) Di Paolo, N.C. and Shayakhmetov, D.M. (2016) *Nat Immunol* 17, 906-13.
- (5) Lamacchia, C. et al. (2013) *Cytokine* 63, 135-44.
- (6) Rider, P. et al. (2013) *Semin Immunol* 25, 430-8.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

www.cellsignal.com