န္နှ NCAM-L1 (D5N9S) Rabbit mAb





Orders:	877-616-CELL (2355) orders@cellsignal.com			
Support:	877-678-TECH (8324)			
Web:	info@cellsignal.com cellsignal.com			

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 220	Source/Isotype: Rabbit IgG	UniProt ID: #P32004	Entrez-Gene Id: 3897		
Product Usage Information		Application Western Blotting Immunoprecipitation Immunofluorescence	(Immunocytochem	istry)		Dilution 1:1000 1:50 1:400		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sensitivity NCAM-L1 (D5N9S) Rabbit mAb recognizes endogenous levels of total NCAM-L1/L1CAM prote						AM protein.		
Source / Purifi	cation		Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human NCAM-L1/L1CAM protein.					
Background		Neural cell adhesion molecule L1 (NCAM-L1/L1CAM) is a single pass transmembrane glycoprotein member of the immunoglobulin superfamily, containing six amino-terminal extracellular Ig-like domains followed by five fibronectin type-III domains (1). NCAM-L1 is mainly expressed in the brain, and plays an important role in the developing nervous system, with involvement in neurite fasciculation and outgrowth, myelination, neuronal migration, and neuronal cell adhesion (2). Mutations in the <i>NCAM-L1</i> gene cause varying degrees of neurological disease including X-linked hydrocephalus, MASA syndrome, spastic paraplegia type 1, and X-linked corpus callosum agenesis, together known as L1 syndrome (3). Apart from the nervous system, NCAM-L1 is overexpressed in many cancers and supports a poor prognosis by facilitating aggressive tumor growth, metastasis, and chemoresistance (4,5).						
Background Re	eferences	1. Moos, M. et al. (1988) <i>Nature</i> 334, 701-3. 2. Hortsch, M. (1996) <i>Neuron</i> 17, 587-93. 3. Christaller, W.A. et al. (2017) <i>Clin Genet</i> 91, 115-120. 4. Kiefel, H. et al. (2012) <i>Cell Adh Migr</i> 6, 374-84. 5. Jo, D.H. et al. (2017) <i>Oncotarget</i> 8, 15441-15452.						
Species Reacti	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	еу	W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry)						
Cross-Reactivit	ty Key	H: Human Mk: Monkey						
Trademarks ar	nd Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		XP is a registered trademark of Cell Signaling Technology, Inc.						
		All other trademarks a more information.	re the property of t	heir respective owners.	Visit cellsignal.com	n/trademarks for		
Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorize the following terms apply to Products provided by CST, its affiliates or its distriterms and conditions that are in addition to, or different from, those contained separately accepted in writing by a legally authorized representative of CST, are force or effect.					es or its distributor ose contained here	s. Any Customer's ein, unless		

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.