



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#9012 Store at -20C

BRCA2 Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M B	Sensitivity: Endogenous	MW (kDa): 380	Source/Isotype: Rabbit	UniProt ID: #P51587	Entrez-Gene Id: 675
---------------------------	-----------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	-------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

BRCA2 Antibody detects endogenous levels of total BRCA2 protein. The antibody does not recognize BRCA1.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acids near the amino terminus of human BRCA2. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The breast cancer susceptibility proteins BRCA1 and BRCA2 are frequently mutated in cases of hereditary breast and ovarian cancers and have roles in multiple processes related to DNA damage, repair, cell cycle progression, transcription, ubiquitination, and apoptosis (1-4). BRCA2 has been shown to be required for localization of Rad51 to sites of double-stranded breaks (DSBs) in DNA, and cells lacking BRCA1 and BRCA2 cannot repair DSBs through the Rad51-dependent process of homologous recombination (HR) (5). Numerous DNA damage-induced phosphorylation sites on BRCA1 have been identified, including Ser988, 1189, 1387, 1423, 1457, 1524, and 1542, and kinases activated in a cell cycle-dependent manner, including Aurora A and CDK2, can also phosphorylate BRCA1 at Ser308 and Ser1497, respectively (6-10). Cell cycle-dependent phosphorylation of BRCA2 at Ser3291 by CDKs has been proposed as a mechanism to switch off HR as cells progress beyond S-phase by blocking the carboxy-terminal Rad51 binding site (11).

Background References

1. Rahman, N. and Stratton, M.R. (1998) *Annu Rev Genet* 32, 95-121.
2. Gayther, S.A. et al. (1999) *Am J Hum Genet* 65, 1021-9.
3. Kerr, P. and Ashworth, A. (2001) *Curr Biol* 11, R668-76.
4. Scully, R. and Livingston, D.M. (2000) *Nature* 408, 429-32.
5. Tutt, A. and Ashworth, A. (2002) *Trends Mol Med* 8, 571-6.
6. Okada, S. and Ouchi, T. (2003) *J Biol Chem* 278, 2015-20.
7. Cortez, D. et al. (1999) *Science* 286, 1162-6.
8. Xu, B. et al. (2002) *Cancer Res* 62, 4588-91.
9. Ouchi, M. et al. (2004) *J Biol Chem* 279, 19643-8.
10. Ruffner, H. et al. (1999) *Mol Cell Biol* 19, 4843-54.
11. Esashi, F. et al. (2005) *Nature* 434, 598-604.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **B:** Bovine

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.