

Dasatinib

✓ 5 mg


Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/27/18

For Research Use Only. Not For Use In Diagnostic Procedures.

Background: Dasatinib is a small molecule inhibitor of both the Src and Bcr-Abl tyrosine kinases, with IC_{50} values of 0.55 and 3.0 nM for the isolated kinases, respectively (1). Imatinib and dasatinib have similar inhibitory mechanisms, but dasatinib displays greater potency in imatinib-resistant cell lines and greater inhibitory effect on platelet derived growth factor (PDGF) (2,3).

Molecular Formula: $C_{22}H_{26}ClN_7O_2S$




Molecular Weight: 488.01 g/mol

Solubility: Soluble in DMSO at 200 mg/mL; very poorly soluble in ethanol and water with maximum solubility in water estimated to be about 10 μ M.

Purity: >99%.

Directions for Use: Dasatinib is supplied as a lyophilized powder. For a 5 mM stock, reconstitute the 5 mg in 2.05 mL DMSO. Working concentrations and length of treatments vary depending on the desired effect, but it is typically used at 50-500 nM for 2-6 hours.



Western blot analysis of extracts from LNCaP cells, untreated (-) or treated with dasatinib (100 nM, 2 hr; +), using Phospho-Src Family (Tyr416) Antibody #2101 (upper) and Src (36D10) Rabbit mAb #2109 (lower).

Storage: Store lyophilized or in solution at -20°C, desiccated. Protect from light. In lyophilized form, the chemical is stable for 24 months. Once in solution, use within 3 months to prevent loss of potency. Aliquot to avoid multiple freeze/thaw cycles.

Background References:

- (1) Lombardo, L.J. et al. (2004) *J Med Chem* 47, 6658-61.
- (2) Copland, M. et al. (2006) *Blood* 107, 4532-9.
- (3) Chen, Z. et al. (2006) *Mol Pharmacol* 69, 1527-33.