#91144

re at -20C	COL1A1 (E8I9Z) Rabbit mAb
Stol	



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 220	Source/Isotype: Rabbit IgG	UniProt ID: #P02452	Entrez-Gene Id: 1277	
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:100		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		COL1A1 (E8I9Z) Rabbit mAb recognizes endogenous levels of total COL1A1 protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Phe1197 of human COL1A1 protein.					
Background		Type 1 collagen is the most abundant collagen in many human tissues, including bone, skin, and tendons. It is a trimeric complex composed of two molecules of COL1A1 (alpha-1 type 1 collagen) and one molecule of COL1A2 (alpha-2 type 1 collagen) (1-3). The expression levels of COL1A1 are regulated by multiple mechanisms, including mRNA stability, translation, and post-translational modification (3-5). Overexpression of COL1A1 has been positively associated with tissue fibrosis disorders, including systemic sclerosis (6), while loss-of-function mutations in the <i>COL1A1</i> gene are a major causative factor for osteogenesis imperfecta (brittle bone disease) (7). Notably, COL1A1 expression levels have also been associated with tumor development in gastric, lung, thyroid, and breast cancers. Research studies suggest that upregulation of COL1A1 can generate a modified extracellular matrix environment that promotes cancer cell survival, proliferation, metastasis, and invasion (8-11).					
Background References		2. Chang, S.W. et al. (2 3. Zhang, Y. and Stefa 4. Parsons, C.J. et al. (2 5. Cai, L. et al. (2010) <i>J</i> 6. Jimenez, S.A. and Sa 7. Forlino, A. et al. (20 8. Li, J. et al. (2016) <i>W</i>	2012) <i>Biophys J</i> 102, novic, B. (2016) <i>Int</i> 2011) <i>J Biol Chem</i> 2 <i>Mol Biol</i> 395, 309-2 aitta, B. (1999) <i>Sprir</i> 11) <i>Nat Rev Endocr</i> <i>prld J Surg Oncol</i> 14 I. (2017) <i>J</i> Cancer Re 2017) <i>Breast Cance</i>	<i> Mol Sci</i> 17, 419. 86, 8609-19. 9 <i>ger Semin Immunopath</i> <i>inol</i> 7, 540-57. , 297. 9s Clin Oncol 143, 1133-4 9r <i>Res</i> 19, 9.	nol 21, 397-414.		
Species Reactiv	vity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).	
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key		W: Western Blotting IP: Immunoprecipitation					
Cross-Reactivity Key		H: Human					
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.					
		All other trademarks a more information.	are the property of	their respective owners.	Visit cellsignal.com	/trademarks for	
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless					

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.