PIAS1 (D33A7) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355) orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M R Mk	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #075925	Entrez-Gene Id: 8554
Product Usage Information		Application Flow Cytometry (Fixed/P	ermeabilized)		Dilution 1:50
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.			
Specificity/Sensitivity		PIAS1 (D33A7) XP [®] Rabbit mAb (PE Conjugate) recognizes endogenous levels of total PIAS1 protein.			
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser550 of human PIAS1 protein.			
Description		This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated PIAS1 (D33A7) XP [®] Rabbit mAb #3550.			
Background		The protein inhibitor of activated Stat (PIAS) proteins, which include PIAS1, PIAS3, PIASx, and PIASy, were originally characterized based on their interaction with the Stat family of transcription factors (1,2). PIAS1, PIAS3, and PIASx interact with and repress Stat1, Stat3, and Stat4, respectively (1-3). Deletion of PIAS1 leads to inhibition of interferon-inducible genes and increased protection against infection (4). The PIAS family contains a conserved RING domain that has been linked to a function as a small ubiquitin-related modifier (SUMO) ligase, coupling the SUMO conjugating enzyme Ubc9 with its substrate proteins (5,6). Numerous studies have now shown that PIAS family members can regulate the activity of transcription factors through distinct mechanisms, including NF-kB (7,8), c-Jun, p53 (5,9), Oct-4 (10), and Smads (11,12). The activity of PIAS1 is regulated by both phosphorylation and arginine methylation. Inflammatory stimuli can induce IKK-mediated phosphorylation of PIAS1 at Ser90, which is required for its activity (13). In addition, PRMT1 induces arginine methylation of PIAS1 at Arg303 following interferon treatment and is associated with its repressive activity on Stat1 (14).			
Background References		1. Liu, B. et al. (1998) <i>Proc Natl Acad Sci USA</i> 95, 10626-31. 2. Chung, C.D. et al. (1997) <i>Science</i> 278, 1803-5. 3. Arora, T. et al. (2003) <i>J Biol Chem</i> 278, 21327-30. 4. Liu, B. et al. (2004) <i>Nat Immunol</i> 5, 891-8. 5. Schmidt, D. and Müller, S. (2002) <i>Proc Natl Acad Sci USA</i> 99, 2872-7. 6. Kotaja, N. et al. (2005) <i>Mol Cell Biol</i> 22, 5222-34. 7. Liu, B. et al. (2005) <i>Mol Cell Biol</i> 25, 1113-23. 8. Tahk, S. et al. (2007) <i>Proc Natl Acad Sci USA</i> 104, 11643-8. 9. Bischof, O. et al. (2006) <i>Mol Cell</i> 22, 783-94. 10. Tolkunova, E. et al. (2007) <i>J Mol Biol</i> 374, 1200-12. 11. Long, J. et al. (2004) <i>Proc Natl Acad Sci USA</i> 101, 99-104. 12. Murdoch, R.N. and Edwards, T. (1992) <i>Biochem Int</i> 28, 1029-37. 13. Liu, B. et al. (2007) <i>Genes Dev</i> 23, 118-32.			

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.