

CPSF6 (E4F5F) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignaling.com

Support: 877-678-TECH (8324)

Web: info@cellsignaling.com
cellsignaling.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R Mk	Endogenous	68	Rabbit IgG	#Q16630	11052

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

CPSF6 (E4F5F) Rabbit mAb recognizes endogenous levels of total CPSF6 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu56 of human CPSF6 protein.

Background

Cleavage and polyadenylation of pre-mRNA is regulated by a core group of proteins called the cleavage and polyadenylation specificity factors (CPSF) (1,2). The CPSF proteins interact with poly(A) polymerase (PAP) to recognize the AAUAAA sequence motif and add poly(A). CPSF can also interact with downstream cleavage factors to precisely cleave the 3' end of pre-mRNA (2,3). CPSF is brought to 3' ends by the carboxy-terminal domain of the Rpb1 subunit of the RNA Polymerase II complex, where it dissociates and initiates polyadenylation (4). CPSF has been shown to have numerous interactions with viral proteins. The influenza NS1 viral protein binds to CPSF4 to prevent 3' end processing of viral RNAs, inhibiting nuclear export (5).

CPSF6 is a 68 kDa member of the CPSF group that has been shown to target the HIV-1 viral capsid (CA) (6). A truncated form of CPSF6, CPSF6-358, localizes exclusively in the cytoplasm and can oligomerize to bind CA, restricting HIV infection and nuclear entry (7,8).

Background References

1. Rügsegger, U. et al. (1996) *J Biol Chem* 271, 6107-13.
2. Murthy, K.G. and Manley, J.L. (1992) *J Biol Chem* 267, 14804-11.
3. Bienroth, S. et al. (1993) *EMBO J* 12, 585-94.
4. Hirose, Y. and Manley, J.L. (1998) *Nature* 395, 93-6.
5. Nemeroff, M.E. et al. (1998) *Mol Cell* 1, 991-1000.
6. Lee, K. et al. (2010) *Cell Host Microbe* 7, 221-33.
7. Lee, K. et al. (2012) *J Virol* 86, 3851-60.
8. Ning, J. et al. (2018) *J Virol* 92, pii: e00368-18. doi: 10.1128/JVI.00368-18.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignaling.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.