Phospho-GSK-3α/β (Ser21/9) Antibody





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk Z	Sensitivity: Endogenous	MW (kDa): 46 GSK-3beta. 51 GSK-3alpha.	Source/Isotype: Rabbit	UniProt ID: #P49840, #P49841	Entrez-Gene Id: 2931, 2932		
Product Usage Information		Application Western Blotting		Dilution 1:1000				
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. 20°C. Do not aliquot the antibody.				cerol. Store at –		
Specificity/Sensitivity		Phospho-GSK-3alpha/beta (Ser21/9) Antibody detects endogenous levels of GSK-3 only when phosphorylated at serine 21 of GSK-3alpha or serine 9 of GSK-3beta.						
Source / Purification		Polyclonal antibodies are produced by immunizing animals with synthetic phosphopeptides corresponding to the sequences of human GSK-3 alpha and beta. Antibodies are purified by protein A and peptide affinity chromatography.						
Background	Glycogen synthase kinase-3 (GSK-3) was initially identified as an enzyme that regulates glycogen synthesis in response to insulin (1). GSK-3 is a ubiquitously expressed serine/threonine protein kina that phosphorylates and inactivates glycogen synthase. GSK-3 is a critical downstream element of t PI3K/Akt cell survival pathway whose activity can be inhibited by Akt-mediated phosphorylation at Ser21 of GSK-3α and Ser9 of GSK-3β (2,3). GSK-3 has been implicated in the regulation of cell fate in <i>Dictyostelium</i> and is a component of the Wnt signaling pathway required for <i>Drosophila, Xenopus,</i> mammalian development (4). GSK-3 has been shown to regulate cyclin D1 proteolysis and subcellul localization (5).							
Background Re	und References 1. Welsh, G.I. et al. (1996) Trends Cell Biol 6, 274-9. 2. Srivastava, A.K. and Pandey, S.K. (1998) Mol Cell Biochem 182, 135-41. 3. Cross, D.A. et al. (1995) Nature 378, 785-9. 4. Nusse, R. (1997) Cell 89, 321-3. 5. Diehl, J.A. et al. (1998) Genes Dev 12, 3499-511. 6. Devi, L. and Ohno, M. (2015) Transl Psychiatry 5, e562.							
Species Reactiv	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot B	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				5% w/v BSA, 1X		
Applications K	ey	W: Western Blotting						
Cross-Reactivit	ty Key	H: Human M: Mouse R: Rat Mk: Monkey Z: Zebrafish						
Trademarks ar	nd Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				Any Customer's , unless		
		Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in						

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products on services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.