

Store at
-20°C

#93659

Total β -Catenin Matched Antibody Pair

Support: +1-978-867-2388 (U.S.)
cellsignal.com/support

Orders: 877-616-2355 (U.S.)
orders@cellsignal.com

Entrez-Gene ID #1499
UniProt ID #P35222

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Isotype
β -Catenin (L54E2) Mouse mAb (BSA and Azide Free)	34396	100 μ g	Mouse IgG1
β -Catenin (D10A8) XP [®] Rabbit mAb (BSA and Azide Free)	84441	100 μ g	Rabbit IgG

Species Reactivity: H, M, Mk


Description: The Total β -Catenin Matched Antibody Pair is ideal for use with immunoassay technologies and high throughput ELISA platforms requiring antibody pairs with specialized or custom antibody labeling. Labels include fluorophores, lanthanides, biotin, and beads. Platforms requiring conjugated Matched Antibody Pairs include MSD, Quanterix Simoa, Alpha Technology (AlphaScreen, AlphaLISA, LANCE, HTRF), and Luminex.

Learn how Matched Antibody Pairs move your projects forward, faster at cst-science.com/matched-antibody-pairs.

Background: β -catenin is a key downstream effector in the Wnt signaling pathway (1). It is implicated in two major biological processes in vertebrates: early embryonic development (2) and tumorigenesis (3). CK1 phosphorylates β -catenin at Ser45. This phosphorylation event primes β -catenin for subsequent phosphorylation by GSK-3 β (4-6). GSK-3 β destabilizes β -catenin by phosphorylating it at Ser33, Ser37, and Thr41 (7). Mutations at these sites result in the stabilization of β -catenin protein levels and have been found in many tumor cell lines (8).

Specificity/Sensitivity: This kit detects proteins from the indicated species, as determined through in-house testing, but may also detect homologous proteins from other species.

Formulation: Supplied in 1X PBS (10 mM Na₂HPO₄, 3 mM KCl, 2 mM KH₂PO₄, and 140 mM NaCl (pH 7.8)). BSA and Azide Free.



Data using Total β -Catenin Matched Antibody Pair #93659 are shown. The two antibodies function as a capture-target-detection sandwich (#34396 and #84441, respectively) to detect endogenous protein expression in treated and untreated cells.

Storage: Store at -20°C. This product will freeze at -20°C so it is recommended to aliquot into single-use vials to avoid multiple freeze/thaw cycles. A slight precipitate may be present and can be dissolved by gently vortexing. This will not interfere with antibody performance.

Please visit cellsignal.com for validation data and a complete listing of recommended companion products.

Directions for Use: Matched Antibody Pairs include capture and detection antibodies to non-overlapping epitopes. Optimal dilutions/concentrations should be determined by the end user.

Background References:

- (1) Cadigan, K.M. and Nusse, R. (1997) *Genes Dev* 11, 3286-3305.
- (2) Wodarz, A. and Nusse, R. (1998) *Annu Rev Cell Dev Biol* 14, 59-88.
- (3) Polakis, P. (1999) *Curr Opin Genet Dev* 9, 15-21.
- (4) Amit, S. et al. (2002) *Genes Dev* 16, 1066-76.
- (5) Liu, C. et al. (2002) *Cell* 108, 837-47.
- (6) Yanagawa, S. et al. (2002) *EMBO J* 21, 1733-42.
- (7) Yost, C. et al. (1996) *Genes Dev* 10, 1443-54.
- (8) Morin, P.J. et al. (1997) *Science* 275, 1787-90.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

cellsignal.com