

#94170
store at +4C**CD44 (IM7) Rat mAb (PE-Cy5[®] Conjugate)**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
FC-FP, FC-L	H M	Endogenous	Rat IgG2b kappa	#P16070	960

Product Usage Information

For optimal flow cytometry results, we recommend 0.125µg of antibody per test.

Application

Flow Cytometry (Fixed/Permeabilized)
Flow Cytometry (Live)

Dilution

1:160
1:160

Storage

Supplied in 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% NaN₃, 0.1% gelatin, pH7.2. This product is stable for 6 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.

Specificity/Sensitivity

CD44 (IM7) Rat mAb (PE-Cy5[®] Conjugate) recognizes endogenous levels of total CD44 protein. This antibody detects an epitope within the extracellular domain and is expected to detect all isoforms of CD44.

Source / Purification

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.

Description

This Cell Signaling Technology antibody is conjugated to PE-Cy5[®] and tested in-house for direct flow cytometric analysis in human and mouse cells.

Background

CD44 is a type I transmembrane glycoprotein that mediates cell-cell and cell-matrix interaction through its affinity for hyaluronic acid (HA) and possibly through other parts of the extracellular matrix (ECM). CD44 is highly polymorphic, possesses a number of alternative splice variants and undergoes extensive post-translational modifications (1,2). Increased surface levels of CD44 are characteristic of T cell activation, and expression of the protein is upregulated during the inflammatory response. Research studies have shown that interactions between CD44 and HER2 are linked to an increase in ovarian carcinoma cell growth (1-3). CD44 interacts with ezrin, radixin, and moesin (ERM), linking the actin cytoskeleton to the plasma membrane and the ECM (4-6). CD44 is constitutively phosphorylated at Ser325 in resting cells. Activation of PKC results in phosphorylation of Ser291, dephosphorylation of Ser325, disassociation of ezrin from CD44, and directional motility (4).

Background References

1. Goodison, S. et al. (1999) *Mol. Pathol.* 52, 189-196.
2. Cichy, J. and Puré, E. (2003) *J. Cell Biol.* 161, 839-843.
3. Bourguignon, L.Y. et al. (1997) *J. Biol. Chem.* 272, 27913-27918.
4. Legg, J.W. et al. (2002) *Nat. Cell Biol.* 4, 399-407.
5. Yonemura, S. et al. (1998) *J. Cell Biol.* 140, 885-895.
6. Tsukita, S. et al. (1994) *J. Cell Biol.* 126, 391-401.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key

FC-FP: Flow Cytometry (Fixed/Permeabilized) **FC-L:** Flow Cytometry (Live)

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Cy and CyDye are registered trademarks of GE Healthcare.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.