ន្ត Rab6 (D37C7) Rabbit mAb





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

_

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 24	Source/Isotype: Rabbit	UniProt ID: #P20340	Entrez-Gene Id: 5870		
Product Usage Information		Application Western Blotting Immunoprecipitation Immunofluorescence	(Immunocytochem	istry)		Dilution 1:1000 1:50 1:400		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	Rab6 (D37C7) Rabbit mAb recognizes endogenous levels of total Rab6 protein.						
Species predict based on 100% homology		Monkey						
Source / Purific	cation	Monoclonal antibody is produced by immunizing animals with a recombinant protein specific to the carboxy terminus of human Rab6 protein.						
Background		Rab6 is a member of the Ras superfamily of small Rab GTPases implicated in endocytosis (1). The three distinct members of the Rab6 subfamily (Rab6A, Rab6A', and Rab6B) are structurally similar but likely exhibit non-overlapping functions (2,3). Rab6 localized to the Golgi (4) regulates retrograde transport of membrane-bound target proteins from the Golgi apparatus to the endoplasmic reticulum (5-7) or from the Golgi to the endosome during exocytotic transport (8). Rab6 interacts with microtubule motor proteins such as rabkinesin-6 (KIF20A) and dynein/dynactin complexes; Rab6-mediated transport requires a functionally intact microtubule system (9,10). Rab6 also regulates cytokinesis and cell cycle check point through interactions with Rab6 effector proteins, including the dynein/dynactin protein DCTN1 and the GTPase activating protein RABGAP1 (11,12).						
Background Re	eferences	 Zerial, M. and McBride, H. (2001) <i>Nat Rev Mol Cell Biol</i> 2, 107-17. Echard, A. et al. (2000) <i>Mol Biol Cell</i> 11, 3819-33. Del Nery, E. et al. (2006) <i>Traffic</i> 7, 394-407. Goud, B. et al. (1994) <i>Ann N Y Acad Sci</i> 733, 340-3. White, J. et al. (1999) <i>J Cell Biol</i> 147, 743-60. Opdam, F.J. et al. (2000) <i>J Cell Sci</i> 113 (Pt 15), 2725-35. Martinez, O. et al. (1997) <i>Proc Natl Acad Sci USA</i> 94, 1828-33. Grigoriev, I. et al. (2007) <i>Dev Cell</i> 13, 305-14. Echard, A. et al. (1998) <i>Science</i> 279, 580-5. Young, J. et al. (2005) <i>Mol Biol Cell</i> 16, 162-77. Hill, E. et al. (2000) <i>EMBO J</i> 19, 5711-9. Miserey-Lenkei, S. et al. (2006) <i>EMBO J</i> 25, 278-89. 						
Species Reactiv	vity	Species reactivity is de	termined by testin	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot B	uffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry)						
Cross-Reactivit	су Кеу	H: Human M: Mouse R: Rat						
Trademarks an	d Patents	Cell Signaling Technol	ogy is a trademark	of Cell Signaling Techno	logy, Inc.			
		All other trademarks a more information.	re the property of	their respective owners.	Visit cellsignal.com	/trademarks for		

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.