

#9638
 Store at -20°C

14-3-3 τ Antibody




Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 04/22/16

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Species Cross-Reactivity*	Molecular Wt.	Source
W, IHC-P Endogenous	H, M, R, Mk	28 kDa	Rabbit**


Background: The 14-3-3 family of proteins plays a key regulatory role in signal transduction, checkpoint control, apoptotic and nutrient-sensing pathways (1,2). 14-3-3 proteins are highly conserved and ubiquitously expressed. There are at least seven isoforms, β , γ , ϵ , ζ , τ and η that have been identified in mammals. The initially described α and δ isoforms are confirmed to be phosphorylated forms of β and ζ , respectively (3). Through their amino-terminal α helical region, 14-3-3 proteins forms homo- or heterodimers that interact with a wide variety of proteins: transcription factors, metabolic enzymes, cytoskeletal proteins, kinases, phosphatases and other signaling molecules (3,4). The interaction of 14-3-3 proteins with their targets is primarily through a phospho-Ser/Thr motif. However, binding to divergent phospho-Ser/Thr motifs, as well as phosphorylation independent interactions have been observed (4). 14-3-3 binding masks specific sequences of the target protein, and therefore, modulates target protein localization, phosphorylation state, stability and molecular interactions (1-4). 14-3-3 proteins may also induce target protein conformational changes which modify target protein function (4,5). Distinct temporal and spatial expression patterns of 14-3-3 isoforms have been observed in development and in acute response to extracellular signals and drugs, suggesting that 14-3-3 isoforms may perform different functions despite their sequence similarities (4). Several studies suggest that 14-3-3 isoforms are differentially regulated in cancer and neurological syndromes (2,3).



Western blot analysis of extracts from HeLa, NIH/3T3 and C6 cells, using 14-3-3 τ Antibody.

Specificity/Sensitivity: 14-3-3 τ Antibody detects endogenous levels of total 14-3-3 τ protein.

Source/Purification: Polyclonal antibodies are produced by immunizing rabbits with a synthetic peptide (KLH-coupled) derived from the sequence of human 14-3-3 τ . Antibodies are purified by protein A and peptide affinity chromatography.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma, using 14-3-3 τ Antibody in the presence of control peptide (left) or antigen-specific peptide (right).

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu\text{g/ml}$ BSA and 50% glycerol. Store at -20°C . Do not aliquot the antibody.

***Species cross-reactivity is determined by western blot.**

****Anti-rabbit secondary antibodies must be used to detect this antibody.**

Recommended Antibody Dilutions:

Western blotting	1:1000
Immunohistochemistry (Paraffin)	1:50
Unmasking buffer:	Citrate
Antibody diluent:	SignalStain [®] Antibody Diluent #8112

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.


Background References:

- (1) Muslin, A.J. and Xing, H. (2000) *Cell. Signal.* 12, 703-709.
- (2) Mackintosh, C. (2004) *Biochem. J.* 381, 329-342.
- (3) Dougherty, M.K. and Morrison, D.K. (2004) *J. Cell Sci.* 117, 1875-1884.
- (4) Yaffe, M.B. (2002) *FEBS Lett.* 513, 53-57.
- (5) Bridges, D. and Moorhead, G.B. (2004) *Sci. STKE* 2004, re10.


IMPORTANT: For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

© 2012 Cell Signaling Technology, Inc.
 Cell Signaling Technology[®] is a trademark of Cell Signaling Technology, Inc.


Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma, using 14-3-3 τ Antibody.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma, showing cytoplasmic and nuclear localization using 14-3-3 τ Antibody.



Immunohistochemical analysis of paraffin-embedded human GIST, using 14-3-3 τ Antibody.