



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#9878

## Phospho-Numb (Ser276) (D5C2) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IF-IC	H	Endogenous	78	Rabbit IgG	#P49757	8650

### Product Usage Information

#### Application

Western Blotting  
Immunofluorescence (Immunocytochemistry)

#### Dilution

1:1000  
1:200

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Phospho-Numb (Ser276) (D5C2) Rabbit mAb recognizes endogenous levels of numb protein only when phosphorylated at Ser276.

### Species predicted to react based on 100% sequence homology

Mouse, Rat, Chicken, Xenopus, Zebrafish, Bovine, Horse

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser276 of human numb protein.

### Background

Numb contains an amino-terminal phosphotyrosine binding (PTB) domain and carboxy-terminal endocytic binding motifs for α-adaptin and EH (Eps15 homology) domain-containing proteins, indicating a role in endocytosis (1,2). There are four mammalian Numb splicing isoforms that are differentially expressed and may have distinct functions (3-5). Numb acts as a negative regulator of Notch signaling by promoting ubiquitination and degradation of Notch (6). The protein is asymmetrically segregated into one daughter cell during cell division, producing two daughter cells with different responses to Notch signaling and different cell fates (7,8). The localization of Numb can also be regulated by G protein-coupled receptor (GPCR) and protein kinase C (PKC) signaling (9).

Numb can be phosphorylated at several sites including Ser7, Ser276, and Ser295. Phosphorylation at these sites regulates asymmetric membrane localization of Numb and integrin endocytosis (10-12).

### Background References

- Berdnik, D. et al. (2002) *Dev. Cell* 3, 221-231.
- Santolini, E. et al. (2000) *J. Cell Biol.* 151, 1345-1352.
- Dho, S.E. et al. (1999) *J. Biol. Chem.* 274, 33097-33104.
- Verdi, J.M. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 10472-10476.
- Toriya, M. et al. (2006) *Dev. Neurosci.* 28, 142-155.
- McGill, M.A. and McGlade, C.J. (2003) *J. Biol. Chem.* 278, 23196-23203.
- Verdi, J.M. et al. (1996) *Curr. Biol.* 6, 1134-1145.
- Reugels, A.M. et al. (2006) *Dev. Dyn.* 235, 934-948.
- Dho, S.E. et al. (2006) *Mol. Biol. Cell* 17, 4142-4155.
- Nishimura, T. and Kaibuchi, K. (2007) *Dev Cell* 13, 15-28.
- Smith, C.A. et al. (2007) *EMBO J* 26, 468-80.
- Wirtz-Peitz, F. et al. (2008) *Cell* 135, 161-73.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)

### Cross-Reactivity Key

**H:** Human

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.