

Store at
-20°C

#99075

CD83 (D8V7V) Rabbit mAb



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H	Endogenous	23-65	Rabbit IgG	#Q01151	9308

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CD83 (D8V7V) Rabbit mAb recognizes endogenous levels of total CD83 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro63 of human CD83 protein.

Background

CD83 is a single-transmembrane protein with a calculated molecular weight (MW) of 23 kDa, but due to heavy and differential glycosylation, its apparent MW ranges from 23 to 70 kDa. CD83 is predominantly expressed on mature dendritic cells (DCs) and has been used as a DC activation/maturation marker as its increased expression is correlated with upregulation of HLA class II antigen expression on DCs (1-3). CD83 is also expressed at a low level on lymphocytes and is upregulated upon lymphocyte activation (4,5). Thymic epithelial cells also express CD83, which is required for normal CD4⁺ T cell development (6,7). CD83 is also expressed as a soluble form (sCD83) that can be found in serum of healthy adults (8). sCD83 has been shown to negatively regulate immune response by lymphocytes (9,10).

Background References

1. Zhou, L.J. et al. (1992) *J Immunol* 149, 735-42.
2. Prechtel, A.T. and Steinkasserer, A. (2007) *Arch Dermatol Res* 299, 59-69.
3. Prazma, C.M. and Tedder, T.F. (2008) *Immunol Lett* 115, 1-8.
4. Prazma, C.M. et al. (2007) *J Immunol* 179, 4550-62.
5. Lüthje, K. et al. (2008) *Int Immunol* 20, 949-60.
6. Fujimoto, Y. et al. (2002) *Cell* 108, 755-67.
7. Breloer, M. and Fleischer, B. (2008) *Trends Immunol* 29, 186-94.
8. Hock, B.D. et al. (2001) *Int Immunol* 13, 959-67.
9. Lin, W. et al. (2017) *Sci Rep* 7, 12895.
10. Horvatinovich, J.M. et al. (2017) *J Immunol* 198, 2286-2301.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.