# Store at -20C

# Translational Control Antibody Sampler Kit



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

1 Kit (6 x 20 microliters)

### For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Phospho-Akt (Ser473) (D9E) XP <sup>®</sup> Rabbit mAb	4060	20 µl	60 kDa	Rabbit IgG
Phospho-p70 S6 Kinase (Thr389) (108D2) Rabbit mAb	9234	20 µl	70, 85 kDa	Rabbit IgG
Phospho-S6 Ribosomal Protein (Ser235/236) (D57.2.2E) XP <sup>®</sup> Rabbit mAb	4858	20 µl	32 kDa	Rabbit IgG
Phospho-eIF2α (Ser51) (D9G8) XP <sup>®</sup> Rabbit mAb	3398	20 µl	38 kDa	Rabbit IgG
Phospho-4E-BP1 (Thr37/46) (236B4) Rabbit mAb	2855	20 µl	15 to 20 kDa	Rabbit IgG
Phospho-eIF4E (Ser209) Antibody	9741	20 µl	25 kDa	Rabbit
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Translational Control Antibody Sampler Kit provides a fast and economical means of evaluating multiple proteins involved in translational control. The kit contains enough primary and secondary antibody to perform two Western blot experiments.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl,  $100 \mu g/ml$  BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Background** 

Key steps in translational control occur at the level of eukaryotic initiation factor 4F (eIF4F) and p70 S6 kinase regulation. eIF4F is a complex whose functions include the recognition of the mRNA 5' cap structure. Several stimuli, such as insulin and various growth and survival factors, regulate the eIF4F complex and p70 S6 kinase primarily by triggering a signaling cascade dependent on sequential activation of PI3K, Akt/PKB and mTOR/FRAP kinases. Akt is activated by phosphorylation within the Cterminus at Ser473 and within the activation loop at Thr308 by phospholipid-dependent kinases. Inactivation in vivo of PI3K by the highly selective inhibitor LY294002 inhibits Akt and downstream elements of this cascade. Direct phosphorylation of mTOR/FRAP at Ser2448 by Akt is a key regulatory event controlling its kinase activity. mTOR/FRAP activity can be effectively blocked by Rapamycin, leading to inactivation of eukaryotic initiation factor 4E binding protein 1 (4E-BP1), an inhibitor of translation initiation, and activation of p70 S6 kinases. Inactivation of 4E-BP1 by sequential phosphorylation causes the release of eIF4E, which, together with eIF4G and other factors, forms a functional eIF4F cap binding complex. p70 S6 kinases phosphorylates the 40S ribosomal subunit protein S6 and stimulates the translation of 5' oligopyrimidine tract containing mRNAs. The Erk pathway is also involved in regulation at this level by regulating the eIF4E kinase, Mnk1, and activating p70 S6 kinase. Tuberin, a product of the tumor supressor gene TSG2, is directly phosphorylated atThr1462 by Akt/PKB. Tuberin inhibits the mammalian target of rapamycin, mTOR, which results in inhibition of p70 S6 kinase and activation of 4E-BP1 and, therefore, inhibition of translation.

### **Background References**

- 1. Gingras, A. et al. (1999) *Annu. Rev. Biochem.* 68, 913-963.
- 2. Gingras, A. C. et al. (2001) *Genes and Develop.* 15, 807-826.
- 3. Dennis, P. B. et al. (1999) Curr. Opin. Genet. Dev. 9, 49-54.
- 4. Volarevic, S. and Thomas, G. (2000) Prog. Nucleic Acid Res. Mol. Biol. 65, 101-127.
- 5. Pyronnet, S. et al. (2000) *Biochem. Pharmacol.* 60, 1237-1243.
- 6. Dever, T.E. (2002) Cell 108, 545-56.
- 7. Goncharova, E.A. et al. (2002) J Biol Chem 277, 30958-67.

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.