



Orders: 877-616-CELL (2355)  
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#9936

## NF- $\kappa$ B Pathway Antibody Sampler Kit

1 Kit (7 x 20 microliters)

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
IKK $\alpha$ (3G12) Mouse mAb	11930	20 $\mu$ l	85 kDa	Mouse IgG1
IKK $\beta$ (D30C6) Rabbit mAb	8943	20 $\mu$ l	87 kDa	Rabbit IgG
Phospho-IKK $\alpha$ / $\beta$ (Ser176/180) (16A6) Rabbit mAb	2697	20 $\mu$ l	85 IKK-alpha 87 IKK-beta kDa	Rabbit IgG
Phospho-NF- $\kappa$ B p65 (Ser536) (93H1) Rabbit mAb	3033	20 $\mu$ l	65 kDa	Rabbit IgG
I $\kappa$ B $\alpha$ (L35A5) Mouse mAb (Amino-terminal Antigen)	4814	20 $\mu$ l	39 kDa	Mouse IgG1
Phospho-I $\kappa$ B $\alpha$ (Ser32) (14D4) Rabbit mAb	2859	20 $\mu$ l	40 kDa	Rabbit IgG
NF- $\kappa$ B p65 (D14E12) XP <sup>®</sup> Rabbit mAb	8242	20 $\mu$ l	65 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 $\mu$ l		Goat
Anti-mouse IgG, HRP-linked Antibody	7076	100 $\mu$ l		Horse

Please visit [cellsignal.com](http://cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

### Description

The NF- $\kappa$ B Pathway Antibody Sampler Kit contains reagents to examine the activation state and total protein levels of key proteins in the NF- $\kappa$ B pathway: IKK $\alpha$ , IKK $\beta$ , NF- $\kappa$ B p65/RelA, and I $\kappa$ B $\alpha$ . The kit contains enough primary and secondary antibodies to perform two Western blot experiments per primary antibody.

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100  $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

### Background

The transcriptional nuclear factor  $\kappa$ B (NF- $\kappa$ B)/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory I $\kappa$ B proteins. Activation occurs via phosphorylation of I $\kappa$ B $\alpha$  at Ser32 and Ser36, resulting in the ubiquitin-mediated proteasome-dependent degradation of I $\kappa$ B $\alpha$  and the release and nuclear translocation of active NF- $\kappa$ B dimers. The regulation of I $\kappa$ B $\beta$  and I $\kappa$ B $\epsilon$  is similar to that of I $\kappa$ B $\alpha$ , however, the phosphorylation and degradation of these proteins occurs with much slower kinetics. Phosphorylation of I $\kappa$ B $\beta$  occurs at Ser/Thr19 and Ser23, while I $\kappa$ B $\epsilon$  can be phosphorylated at Ser18 and Ser22. The key regulatory step in this pathway involves activation of a high molecular weight I $\kappa$ B kinase (IKK) complex, consisting of three tightly associated IKK subunits. IKK $\alpha$  and IKK $\beta$  serve as the catalytic subunits of the kinase. Activation of IKK depends on phosphorylation at Ser177 and Ser181 in the activation loop of IKK $\beta$  (176 and 180 in IKK $\alpha$ ). NF- $\kappa$ B-inducing kinase (NIK), TANK-binding kinase 1 (TBK1), and its homolog IKK $\epsilon$  (IKKi), phosphorylate and activate IKK $\alpha$  and IKK $\beta$ .

The NF- $\kappa$ B family of transcription factors is comprised of five proteins in mammals, p65/RelA, c-Rel, RelB, NF- $\kappa$ B1 (p105/p50) and NF- $\kappa$ B2 (p100/p52). p105 and p100 are proteolytically processed to produce p50 and p52, respectively. The 50 kDa active form is produced through proteolytic processing following IKK-mediated phosphorylation of p105 at multiple sites (Ser922, 924, 928 and 933), while p100's processing to p52 is induced by phosphorylation of Ser864 and Ser868. The p50 and p52 products form dimeric complexes with Rel proteins, which are then able to bind DNA and regulate transcription. Phosphorylation of p65/RelA at Ser276 by PKA C and MSK1 enhances transcriptional activity. p65 phosphorylation at Ser536 regulates activation, nuclear localization, protein-protein interactions, and transcriptional activity. PMA-induced NF- $\kappa$ B transcriptional activity is dependent on the region of p65 containing the potential phosphorylation sites Ser457, Thr458, Thr464 and Ser468. Phosphorylation of Ser468 by GSK-3 $\beta$  inhibits basal p65 activity.

### Background References

1. Yamamoto, Y. and Gaynor, R.B. (2004) *Trends Biochem. Sci.* 29, 72-79.
2. Ghosh, S. and Karin, M. (2002) *Cell* 109, S81-S96.
3. Viatour, P. et al. (2005) *Trends Biochem. Sci.* 30, 43-52.
4. Ho, C. et al. (2016) *PLOS One* 10,1-22.
5. Beyaz S. et al. (2016) *Nature* 531, 53-58.

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.