

#9968 Store at -20C

Rho-GTPase Antibody Sampler Kit



1 Kit (6 x 20 microliters)

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Cdc42 (11A11) Rabbit mAb	2466	20 µl	21 kDa	Rabbit
Phospho-Rac1/cdc42 (Ser71) Antibody	2461	20 µl	28 kDa	Rabbit
RhoA (67B9) Rabbit mAb	2117	20 µl	21 kDa	Rabbit IgG
RhoB Antibody	2098	20 µl	21 kDa	Rabbit
RhoC (D40E4) Rabbit mAb	3430	20 µl	21 kDa	Rabbit IgG
Rac1/2/3 Antibody	2465	20 µl	21 kDa	Rabbit
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

Please visit cellsignal.com for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

Description

The Rho-GTPase Antibody Sampler Kit contains reagents to examine aspects of cell migration, adhesion, proliferation and differentiation in cells. This kit includes enough primary and secondary antibodies to perform two Western blot experiments per each primary antibody.

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Background

The Rho family of small GTPases, including Rho, Rac, and cdc42, act as molecular switches to regulate processes such as cell migration, adhesion, proliferation and differentiation (1). RhoA, RhoB and RhoC are all highly homologous but appear to have divergent biological functions. The best characterized of these proteins, RhoA, regulates actin contractility, cytokinesis, focal adhesion assembly and cell polarity (2-5). Mammalian Rac exists as three isoforms (Rac1, Rac2 and Rac3) that show high sequence similarity. Well-characterized Rac1 and cdc42 are ubiquitously expressed and play key signaling roles in cytoskeletal reorganization, membrane trafficking, transcriptional regulation, and cell growth and development (6). Phosphorylation of Rac1 at a putative Akt site (Ser71) may limit Rac1 activity through inhibition of GTP binding (7). Rac2 is expressed in cells of hematopoietic origin, while Rac3 is highly expressed in brain and in many other tissues. The Vav family of guanine-nucleotide exchange factors mediates activation of Rho/Rac family small GTPases (8). Negative regulation of Rho-activity members of the p190 RhoGAP family (p190-A and p190-B) may be controlled by Src phosphorylation of Tyr residues, activating the p190 GAP domain (8-10). Furthermore, Rho GDP dissociation inhibitor (RhoGDI) associates with Rho/Rac to negatively regulate nucleotide exchange membrane localization (11).

Background References

1. DerMardirossian, C. and Bokoch, G.M. (2005) *Trends Cell Biol.* 15, 356-363.
2. Bi, D. et al. (2005) *Circ. Res.* 96, 890-897.
3. Kimura, K. et al. (2000) *J. Biol. Chem.* 275, 17233-17236.
4. Barry, S.T. and Critchley, D.R. (1994) *J. Cell Sci.* 107 (Pt 7), 2033-2045.
5. Van Keymeulen, A. et al. (2006) *J. Cell Biol.* 174, 437-445.
6. Wennerberg, K. and Der, C.J. (2004) *J. Cell Sci.* 117, 1301-1312.
7. Kwon, T. et al. (2000) *J. Biol. Chem.* 275, 423-428.
8. Sordella, R. et al. (2003) *Cell* 113, 147-158.
9. Chang, J.H. et al. (1995) *J. Cell Biol.* 130, 355-368.
10. Roof, R.W. et al. (1998) *Mol. Cell Biol.* 18, 7052-7063.
11. Dovas, A. and Couchman, J.R. (2005) *Biochem. J.* 390, 1-9.

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.